arXiv:0905.2017v1 [hep-ph] 13 May 2009

Light Scalar Mesons in Photon-Photon Collisions

N.N. Achasov* and G.N. Shestakov†

Laboratory of Theoretical Physics, S.L. Sobolev Institute for Mathematics, 630090, Novosibirsk, Russia

(Dated: May 13, 2009)

The light scalar mesons, discovered forty years ago, became a challenge for the naive quark-antiquark model from the outset. At present the nontrivial nature of these states is no longer denied practically anybody. We review a substantial contribution of the two-photon physics to understanding the nature of the light scalar mesons and new results for their two-photon production mechanisms, based on the analysis of current experimental data.

PACS numbers: 12.39.-x, 13.40.-f, 13.60.Le, 13.75.Lb

Contents

I. Introduction 1

II. Special Place of the Light Scalar Mesons in the Hadron World, Evidences of Their Four-Quark Structure 2

III. Light Scalars in the Light of Two-Photon Collisions 4
- A. History of Investigations 4
- B. Current Experimental Situation 7
- C. Dynamics of the Reactions $\gamma\gamma\to\pi\pi$ 8
  - 1. Born Contributions and Angular Distributions 8
  - 2. Production Mechanisms of Scalar Resonances 9
- D. Analysis of the New High-Statistics Belle Data 10
- E. Conclusion: Two-Photon Decays on the Nature of the $\sigma(600)$ and $f_0(980)$ mesons 17

IV. Future Trends 18
- A. The $a_0(980)$ Meson in the Reaction $\gamma\gamma\to\pi^0\eta$ 18
- B. The $f_0(980)$ and $a_0(980)$ Mesons near the $\gamma\gamma\to K\bar{K}$ Reaction Thresholds 18

Acknowledgments 19

V. Appendix 19
- A. $\gamma\gamma\to\pi\pi$ 19
- B. $\gamma\gamma\to\pi^0\eta$ 21
- C. $\gamma\gamma\to K\bar{K}$ 22

References 23

I. INTRODUCTION

The scalar channels in the region up to 1 GeV became a stumbling block of QCD because both perturbation theory and sum rules do not work in these channels.[1] At the same time the question on the nature of the light scalar mesons is major for understanding the mechanism of the chiral symmetry realization, arising from the confinement, and hence for understanding the confinement itself.

Hunting the light σ and κ mesons had begun in the sixties already and a preliminary information on the light scalar mesons in Particle Data Group (PDG) Reviews had appeared at that time (see, for example, Rosenfeld *et al.*, 1965, 1967; Barash-Schmidt *et al.*, 1969). The theoretical ground for a search for scalar mesons was the linear σ model (LSM) (Gell-Mann and Levy, 1960; Gell-Mann, 1964; Levy, 1967), which takes into account spontaneous breaking of chiral symmetry and contains pseudoscalar mesons as Goldstone bosons. The surprising thing is that after ten years it has been made clear that LSM could be the low energy realization of QCD. At the end of the sixties and at the beginning of the seventies (Rosenfeld *et al.*, 1967; Barash-Schmidt *et al.*, 1969 ; Rittenberg *et al.*, 1971; Lasinski *et al.*, 1973) there were discovered the narrow light scalar resonances, the isovector $a_0(980)$ and isoscalar $f_0(980)$.[2]

As for the σ and κ mesons, long-standing unsuccessful attempts to prove their existence in a conclusive way entailed general disappointment and an information on

---

*Electronic address: `achasov@math.nsc.ru`

†Electronic address: `shestako@math.nsc.ru`

[1] The point is that, in contrast to classic vector channels, in this region there are not solitary resonances, i.e., scalar resonances, which are not accompanied by a large inseparable from resonance background. Particularly, in the case of the solitary $a_0(980)$ and $f_0(980)$ resonances, the resonance peaks in the $\phi\to\gamma a_0(980)\to\gamma\pi\eta$ and $\phi\to\gamma f_0(980)\to\gamma\pi\pi$ decays would be not observed at all because the differential probabilities of these decays vanish proportionally cubic function of a photon energy in a soft photon region for gauge invariance (Achasov and Ivanchenko, 1989; Achasov and Gubin, 2001; Achasov, 2003), see next Section. The principal role of the chiral background in the fate of the $\sigma(600)$ resonance was demonstrated in the linear σ model (Achasov and Shestakov, 1994a, 2007). The solitary resonance approximation is nothing more than an academic exercise in the light scalar meson case.

[2] In 1977 Jaffe noted that in the MIT bag model, which incorporates confinement phenomenologically, there are light four-quark scalar states (Jaffe, 1977). He suggested also that $a_0(980)$ and $f_0(980)$ might be these states with symbolic structures: $a_0^+(980)=ud\bar{s}\bar{s}$, $a_0^0(980)=(us\bar{u}\bar{s}-ds\bar{d}\bar{s})/\sqrt{2}$, $a_0^-(980)=d\bar{u}s\bar{s}$, and $f_0(980)=(us\bar{u}\bar{s}+ds\bar{d}\bar{s})/\sqrt{2}$. From that time $a_0(980)$ and $f_0(980)$ resonances came into beloved children of the light quark spectroscopy.

these states disappeared from PDG Reviews. One of principal reasons against the σ and κ mesons was the fact that both $\pi\pi$ and πK scattering phase shifts do not pass over 90^0 at putative resonance masses. Nevertheless, experimental and theoretical investigation of processes, in which the σ and κ states could reveal themselves, had been continued.

Situation changes when we showed that in LSM there is a negative background phase in the $\pi\pi \to \pi\pi$ scattering amplitude (Achasov and Shestakov, 1994a) which hides the σ meson with the result that the $\pi\pi$ scattering phase shift does not pass over 90^0 at putative resonance masses. It has been made clear that shielding wide lightest scalar mesons in chiral dynamics is very natural. This idea was picked up and triggered new wave of theoretical and experimental searches for the σ and κ mesons (see, for example, Sannino and Schechter, 1995; Törnqvist, 1995; Ishida, Ishida, Takahashi, Ishida, Takamatsu, and Tsuru, 1996; Harada, Sannino, and Schechter, 1996; Black, Fariborz, Sannino, and Schechter, 1998, 1999; Ishida, 2000). As a result the light σ resonance, since 1996, and the light κ resonance, since 2004, appeared in PDG Reviews.

By now there is an impressive amount of data about the light scalar mesons (Amsler *et al.*, 2008; Spanier, Törnqvist, and Amsler, 2008). The nontrivial nature of these states is no longer denied practically anybody. In particular, there exist numerous evidences in favour of their four-quark structure. These evidences are widely covered in the literature (see, for example, Montanet, 1983, Achasov, Devyanin, and Shestakov, 1984a; Achasov and Ivanchenko, 1989; Achasov, 1990, 1998, 1999, 2001, 2003a, 2003b, 2003c, 2004a, 2004b, 2006, 2008a, 2008b; Achasov and Shestakov, 1991, 1999; Delbourgo and Scadron, 1998; Godfrey and Napolitano, 1999; Tuan, 2001, 2003; Close and Törnqvist, 2002; Alford and Jaffe, 2003; Jaffe and Wilczek, 2003; Amsler and Törnqvist, 2004; Maiani, Polosa, Piccinini, and Riquer, 2004; Jaffe, 2005, 2006; Kalashnikova, Kudryavtsev, Nefediev, Hanhart, and Haidenbauer, 2005; Achasov, Kiselev, and Shestakov, 2006, 2008; Caprini, Colangelo, and Leutwyler, 2006; Bugg, 2006; Fariborz, Jora, and Schechter, 2006, 2008a, 2008b, 2009; Narison, 2006, 2008; Törnqvist, 2007; Klempt and Zaitsev, 2007; Maiani, Polosa, and Riquer, 2007; Pennington, 2007a, 2007b; van Beveren and Rupp, 2007; Bystritsky, Volkov, Kuraev, Bartoš, and M. Sečanský, 2008; Leutwyler, 2008; 't Hooft, Isidori, Maiani, Polosa, and Riquer 2008; Ivashin, and Korchin, 2008; Ebert, D., R.N. Faustov, and V.O. Galkin, 2009). They are presented also in Sections II and III.

One of them is the suppression of the $a_0(980)$ and $f_0(980)$ resonances in the $\gamma\gamma \to \pi^0\eta$ and $\gamma\gamma \to \pi\pi$ reactions, respectively, predicted in 1982 (Achasov, Devyanin, and Shestakov, 1982a, 1982b) and confirmed by experiment (Amsler *et al.*, 2008). The elucidations of the mechanisms of the $\sigma(600)$, $f_0(980)$, and $a_0(980)$ resonance production in the $\gamma\gamma$ collision and their quark structure are intimately related. That is why the studies of the two-photon processes are the important part of the light scalar meson physics.

Recently a qualitative leap had place in the experimental investigations of the $\gamma\gamma \to \pi\pi$ processes (Mori *et al.*, 2003; 2007a; 2007b; Uehara *et al.*, 2008) that proved the theoretical expectations based on the four-quark nature of the light scalar mesons (Achasov, Devyanin, and Shestakov, 1982a, 1982b). The Belle Collaboration published the data on the cross sections for the $\gamma\gamma \to \pi^+\pi^-$ (Mori *et al.*, 2007a, 2007b) and $\gamma\gamma \to \pi^0\pi^0$ (Uehara *et al.*, 2008) reactions, statistics of which are hundreds of times as large as statistics of all previous data. The Belle Collaboration observed for the first time the clear signals of the $f_0(980)$ resonance in the both charge channels. The previous indications for the $f_0(980)$ production in the $\gamma\gamma$ collisions (Marsiske *et al.*,1990; Boyer *et al.*, 1990; Oest *et al.*, 1990; Behrend *et al.*, 1992; Bienlein *et al.*, 1992) were rather indefinite.

In the given paper there are presented the results of the investigation the mechanisms of the $\gamma\gamma \to \pi^+\pi^-$ and $\gamma\gamma \to \pi^0\pi^0$ reactions (see Section III) based on the analysis (Achasov and Shestakov, 2008a, 2008b) of the data Belle (Mori *et al.*, 2007b; Uehara *et al.*, 2008) and our previous investigations of the scalar meson physics in the $\gamma\gamma$-collisions (Achasov, Devyanin, and Shestakov, 1982a, 1982b, 1985; Achasov and Shestakov, 1988, 1992, 1994b, 2005, 2007). We also briefly (sometimes critical) survey analyses of other authors.

The joint analysis of the Belle high-statistics data on the $\gamma\gamma \to \pi^+\pi^-$ and $\gamma\gamma \to \pi^0\pi^0$ reactions is presented and the principal dynamical mechanisms of these processes are elucidated in the energy region up to 1.5 GeV. It is shown that the direct coupling constants of the $\sigma(600)$, $f_0(980)$, and $a_0(980)$resonances with the $\gamma\gamma$ system are small with the result that their decays in the two photon are the four-quark transitions caused by the rescatterings $\sigma \to \pi^+\pi^- \to \gamma\gamma$, $f_0(980) \to K^+K^- \to \gamma\gamma$ and $a_0(980) \to K^+K^- \to \gamma\gamma$ in contrast to the two-photon decays of the classic P wave tensor $q\bar{q}$ mesons $a_2(1320)$, $f_2(1270)$ and $f_2'(1525)$, which are caused by the direct two-quark transitions $q\bar{q} \to \gamma\gamma$ in the main. It is obtained the two-photon widths averaged over resonance mass distributions $\langle\Gamma_{f_0\to\gamma\gamma}\rangle_{\pi\pi} \approx 0.19$ keV, $\langle\Gamma_{a_0\to\gamma\gamma}\rangle_{\pi\eta} \approx 0.125$ keV and $\langle\Gamma_{\sigma\to\gamma\gamma}\rangle_{\pi\pi} \approx 0.45$ keV. In addition, the $f_2(1270) \to \gamma\gamma$ width is obtained $\Gamma_{f_2\to\gamma\gamma}(m^2_{f_2}) \approx 3.8$ keV.

We attend also to the additional possibilities of the investigation of the $a_0(980)$ and $f_0(980)$ resonances in the $\gamma\gamma \to \pi^0\eta$ and $\gamma\gamma \to K\bar{K}$ reactions, which are as yet little studied experimentally (see Section IV).

II. SPECIAL PLACE OF THE LIGHT SCALAR MESONS IN THE HADRON WORLD, EVIDENCES OF THEIR FOUR-QUARK STRUCTURE

Even a cursory examination of PDG Reviews gives an idea of the four-quark structure of the light scalar meson nonet [3], $\sigma(600)$, $\kappa(800)$, $a_0(980)$, and $f_0(980)$,

$$\begin{array}{ccccc} a_0^- & & a_0^0/f_0 & & a_0^+ \\ & \{\kappa\} & & \{\kappa\} & \\ & & \sigma & & \end{array} \qquad (1)$$

inverted (Achasov and Shestakov, 1999) in comparison with the classical P wave $q\bar{q}$ tensor meson nonet $f_2(1270)$, $a_2(1320)$, $K_2^*(1420)$, and $f_2'(1525)$

$$\begin{array}{ccccc} & & f_2' & & \\ & \{K_2^*\} & & \{K_2^*\} & \\ a_2^- & & a_2^0/f_2 & & a_2^+ \end{array}\ , \qquad (2)$$

or also in comparison with the classical S wave vector meson nonet $\rho(770)$, $\omega(782)$, $K^*(892)$, and $\phi(1020)$. [4] In the naive quark model such a nonet cannot be understood as the P wave $q\bar{q}$ nonet, but it can be easy understood as the S wave $q^2\bar{q}^2$ nonet, where $\sigma(600)$ has no strange quarks, $\kappa(800)$ has the s quark, $a_0(980)$ and $f_0(980)$ have the $s\bar{s}$ pair.

The scalar mesons $a_0(980)$ and $f_0(980)$, discovered about forty years ago, became the hard problem for the naive $q\bar{q}$ model from the outset. [5] Really, on the one hand the almost exact degeneration of the masses of the isovector $a_0(980)$ and isoscalar $f_0(980)$ states revealed seemingly the structure $a_0^+(980) = u\bar{d}$, $a_0^0(980) = (u\bar{u} - d\bar{d})/\sqrt{2}$, $a_0^-(980) = d\bar{u}$ and $f_0(980) = (u\bar{u} + d\bar{d})/\sqrt{2}$ similar to the structure of the vector ρ and ω or tensor $a_2(1320)$ and $f_2(1270)$ mesons, but on the other hand, the strong coupling of the $f_0(980)$ with the $K\bar{K}$ channel as if suggested a considerable part of the strange pair $s\bar{s}$ in the wave function of the $f_0(980)$.

At the beginning of eighty it was demonstrated in a series of papers (Achasov, Devyanin, and Shestakov, 1980a, 1980b, 1981a, 1981b, 1981c, 1984a, 1984b) that data on the $f_0(980)$ and $a_0(980)$ resonances, available at that time, can be interpreted in favour of the $q^2\bar{q}^2$ model, i.e., can be explained by using coupling constants of the $f_0(980)$ and $a_0(980)$ states with pseudoscalar mesons superallowed by the Okubo-Zweig-Iizuka (OZI) rule as it is predicted by the $q^2\bar{q}^2$ model. In particular, in these papers there were obtained and specified formulae for scalar resonance propagators with taking into account corrections for finite width in case of strong coupling with two-particle decay channels. Late on, these formulae were used in fitting data of a series of experiments on the $f_0(980)$ and $a_0(980)$ resonance production (see, for example, Achasov *et al.*, 1998a, 1998b, 2000a, 2000b; Akhmetshin *et al.*, 1999a, 1999b; Aloisio *et al.*, 2000a, 2000b; Dubrovin, 2003; Ambrosino *et al.*, 2006, 2007a, 2007b; Bonvicini *et al.*, 2007; Cavoto, 2007; Mori *et al.*, 2007a, 2007b; Uehara *et al.*, 2008; Bossi *et al.*, 2008). Recently, it was shown that the above scalar resonance propagators satisfy the Källen-Lehmann representation in the domain of coupling constants usually used (Achasov and Kiselev, 2004).

At the end of eighties it was shown that the study of the radiative decays $\phi \to \gamma a_0 \to \gamma\pi\eta$ and $\phi \to \gamma f_0 \to \gamma\pi\pi$ can shed light on the problem of $a_0(980)$ and $f_0(980)$ mesons (Achasov and Ivanchenko, 1989). Over the next ten years before experiments (1998) the question was considered from different points of viev (Bramon, Grau, and Pancheri, 1992; Close, Isgur, and Kumano, 1993; Lucio and Napsuciale, 1994; Achasov, 1995; Achasov and Gubin, 1997, 1998; Achasov, Gubin, and Shevchenko, 1997; Achasov, Gubin, and Solodov, 1997).

Now these decays have been studied not only theoretically but also experimentally with the help of the SND (Achasov *et al.*, 1998a, 1998b, 2000a, 2000b) and CMD-2 (Akhmetshin *et al.*, 1999a, 1999b) detectors at Budker Institute of Nuclear Physics in Novosibirsk and the KLOE detector at the DAΦNE ϕ-factory in Frascati (Aloisio *et al.*, 2002a, 2002b; Ambrosino *et al.*, 2006, 2007a, 2007b, 2009a, 2009b; Bini, 2008; Bossi *et al.*, 2008). These experimental data called into being a series of investigations (Achasov and Gubin, 2001, Achasov 2003a, 2003b, Achasov and Kiselev 2003, 2006, 2007a) in which evidences for the four-quark nature of the $f_0(980)$ and $a_0(980)$ states were obtained. Note the clear qualitative one. The isovector $a_0(980)$ resonance is produced in the radiative ϕ meson decay as intensively as the isoscalar $\eta'(958)$ meson containing $\approx 66\%$ of $s\bar{s}$, responsible for the $\phi \approx s\bar{s} \to \gamma s\bar{s} \to \gamma\eta'(958)$ decay. In the two-quark model, $a_0^0(980) = (u\bar{u} - d\bar{d})/\sqrt{2}$, the $\phi \approx s\bar{s} \to \gamma a_0(980)$ decay should be suppressed by the OZI rule. So, experiment, probably, indicates for the presence of the $s\bar{s}$ pair in the isovector $a_0(980)$ state, i.e., for its four-quark nature.

---

[3] To be on the safe side, notice that the linear σ model does not contradict to non-$q\bar{q}$ nature of the low lying scalars because Quantum Fields can contain different virtual particles in different regions of virtuality.

[4] In Eqs. (1) and (2) the mass and isotopic spin third component of states increase bottom-up and from left to right, respectively.

[5] Note here a series of important experiments of seventies in which the $f_0(980)$ and $a_0(980)$ resonances were investigated, Flatte *et al.* (1972), Protopopescu *et al.* (1973), Hyams *et al.* (1973), Grayer *et al.* (1974), Hyams *et al.* (1975), Gay *et al.* (1976), as well as a few theoretical analyses of scalar meson properties relevant to this period, Morgan (1974), Flatte (1976), Martin, Ozmutlu, and Squires (1977), Petersen (1977), Jaffe (1977), Estabrooks (1979), Achasov, Devyanin, and Shestakov (1979). In the last-named paper there was theoretically discovered the fine threshold phenomenon of the $a_0(980) - f_0(980)$ mixing which breaks the isotopic invariance (see also Achasov, Devyanin, and Shestakov, 1981b). Now a rebirth of interest in the $a_0(980) - f_0(980)$ mixing takes place and there appear new suggestions on search for this phenomenon (see, for example, Achasov and Shestakov, 2004a, 2004b; Wu, Zhao, and Zou, 2007; Wu and Zou, 2008; and references in these papers) as well as the first indication for its manifestation in the $f_1(1285) \to \pi^+\pi^-\pi^0$ decay, which was measured with the help of the VES detecor in Protvino (Dorofeev *et al.*, 2007, 2008).



Figure 1 The K^+K^- loop model.



Figure 2 The left and right plots illustrate the fit to the KLOE data for the $\pi^0\eta$ and $\pi^0\pi^0$ mass spectra in the $\phi \to \gamma\pi^0\eta$ (Aloisio at al., 2002a) and $\phi \to \gamma\pi^0\pi^0$ (Aloisio at al., 2002b) decays, respectively. See for details Achasov and Kiselev, 2003, 2006, 2007a.

When basing the experimental investigations (Achasov and Inanchenko, 1989), it was suggested the kaon loop model $\phi \to K^+K^- \to \gamma a_0(980) \to \gamma\pi^0\eta$ and $\phi \to K^+K^- \to \gamma f_0(980) \to \gamma\pi\pi$, see Fig. 1. This model is used in the data treatment and is ratified by experiment, see Fig. 2, (Achasov *et al.*, 1998a, 1998b, 2000a, 2000b; Akhmetshin *et al.*, 1999a, 1999b; Aloisio at al., 2002a, 2002b; Ambrosino *et al.*, 2006, 2007a, 2007b, 2009a, 2009b; Bossi *et al.*, 2008; Di Micco, 2008; Shekhovtsova, Venanzoni, and Pancheri, 2009).

The kaon loop model includes the new nontrivial threshold phenomenon, see Fig. 3. To describe the experimental mass distributions $dBR(\phi \to \gamma R \gamma ab;\, m)/dm \sim |g(m)|^2\omega(m)$ [6] the function $|g(m)|^2$ [7] should be smooth at $m \leq 0.99$ GeV. But gauge invariance requires that $g(m)$ is proportional to the photon energy $\omega(m)$. Stopping the impetuous increase of the function $(\omega(m))^3$ at $\omega(990\,\text{MeV})$=29 MeV is the crucial point in the data description. The K^+K^- loop model solves this problem in the elegant way, see Fig. 3, (Achasov and Gubin, 2001; Achasov, 2001, 2003a, 2003b). In truth this means that $a_0(980)$ and $f_0(980)$ resonances are seen in the radiative decays of ϕ meson owing to the K^+K^- intermediate state, otherwise the maxima in the spectra would be shifted to 900 MeV. So the mechanism of the $a_0(980)$ and $f_0(980)$ mesons production in the ϕ radiative decays is established at a physical level of proof at least.



Figure 3 A new threshold phenomenon in $\phi \to K^+K^- \to \gamma R$ decays. The universal in the K^+K^- loop model function $|g(m)|^2 = |g_R(m)/g_{RK^+K^-}|^2$ is drawn with the solid line. The contributions of the imaginary and real parts of $g(m)$ are drawn with the dashed and dotted lines, respectively.

Both real and imaginary parts of the $\phi \to \gamma R$ amplitude are caused by the K^+K^- intermediate state. The imaginary part is caused by the real K^+K^- intermediate state while the real part is caused by the virtual compact K^+K^- intermediate state, i.e., we are dealing here with the four-quark transition (Achasov, 2001, 2003a, 2003b). Needless to say, radiative four-quark transitions can happen between two $q\bar{q}$ states as well as between $q\bar{q}$ and $q^2\bar{q}^2$ states but their intensities depend strongly on a type of the transitions. A radiative four-quark transition between two $q\bar{q}$ states requires creation and annihilation of an additional $q\bar{q}$ pair, i.e., such a transition is forbidden according to the OZI rule, while a radiative four-quark transition between $q\bar{q}$ and $q^2\bar{q}^2$ states requires only creation of an additional $q\bar{q}$ pair, i.e., such a transition is allowed according to the OZI rule. The consideration of this question from the large N_C expansion standpoint supports a suppression of a radiative four-quark transition between two $q\bar{q}$ states in comparison with a radiative four-quark transition between $q\bar{q}$ and $q^2\bar{q}^2$ states. So, both intensity and mechanism of the $a_0(980)$ and $f_0(980)$ production in the radiative decays of the $\phi(1020)$ meson indicate for their four-quark nature.

Note also that the absence of the decays $J/\psi \to \gamma f_0(980)$, $J/\psi \to a_0(980)\rho$, $J/\psi \to f_0(980)\omega$ against a background of the rather intensive decays into the corresponding classical P wave tensor $q\bar{q}$ resonances $J/\psi \to \gamma f_2(1270)$ (or even $J/\psi \to \gamma f_2'(1525)$), $J/\psi \to a_2(1320)\rho$, $J/\psi \to f_2(1270)\omega$ intrigues against the P wave $q\bar{q}$ structure of the $a_0(980)$ and $f_0(980)$ states (Achasov, 1998, 2002, 2003c).

III. LIGHT SCALARS IN THE LIGHT OF TWO-PHOTON COLLISIONS

A. History of Investigations

Experimental investigations of light scalar mesons in the $\gamma\gamma \to \pi^+\pi^-$, $\gamma\gamma \to \pi^0\pi^0$ and $\gamma\gamma \to \pi^0\eta$ reactions with the e^+e^--colliders began in eighties and have continued up to now. In first decade many groups, DM1, DM1/2, PLUTO, TASSO, CELLO, JADE, Crystal Ball, MARK II, DELCO, and TPC/2γ, took part in that. Only Crystal Ball and JADE could studied the $\pi^0\pi^0$ and $\pi^0\eta$ channels, the others (and JADE) the $\pi^+\pi^-$ channel. For those who wish to read more widely in the contribution

---

[6] Here m is the invariant mass of the ab-state, $R = a_0(980)$ or $f_0(980)$, $ab = \pi^0\eta$ or $\pi^0\pi^0$.

[7] The function $g(m)$ describes the $\phi \to \gamma[a_0(m)/f_0(m)]$ transition vertex.

Table I First conclusions on the $f_0(980)$ production in $\gamma\gamma\to\pi\pi$ (see Field, 1981; Higler, 1981; Wedemeyer, 1981).

Experiments	Conclusions
Crystal Ball	No significant $f_0(980)$
CELLO	Hint of $f_0(980)$
JADE	No evidence for $f_0(980)$
TASSO	Good fit to data book values for $f_2(1270)$ includes $f_0(980)$ ($3\,\sigma$ effect)
MARK II	No significant $f_0(980)$ signal

of this impressive period in light scalar meson physics one can recommend the following reviews and papers: Field, 1981; Higler, 1981; Wedemeyer, 1981; Achasov, Devyanin, and Shestakov, 1982a, 1982b; Edwards *at al.*, 1982; Olsson, 1983; Mennessier, 1983; Kolanoski, 1984, 1986, 1988; Cordier, 1984; Erne, 1984; Barnes, 1985, 1992; Kaloshin and Serebryakov, 1986; Antreasyan *at al.*, 1986; Johnson, 1986; Poppe, 1986; Berger and Wagner, 1987; Kolanoski and Zerwas, 1987; Morgan and Pennington, 1987, 1988; Achasov and Shestakov, 1985, 1988, 1991; Chanowitz, 1988; Hikasa *et al.*, 1992; Morgan, Pennington, and Whalley, 1994.

First results on the $f_0(980)$ resonance production are collected in Tables I and II.

It is reasonable that first conclusions had a qualitative character and data on the $f_0(980)\to\gamma\gamma$ decay width had large errors or were upper bounds. Note as a guide that the TASSO and Crystal Ball results, see Table II, based on the integral luminosity equals to $9.24\,\text{pb}^{-1}$ and $21\,\text{pb}^{-1}$ respectively.

As for the $a_0(980)$ resonance, it was observed in the $\gamma\gamma\to\pi^0\eta$ reaction only in two experiments. The Crystal Ball group (Antreasyan *at al.*, 1986) collected during two years the integral luminosity of $110\,\text{pb}^{-1}$, selected at that 336 events relevant to the $\gamma\gamma\to\pi^0\eta$ reaction in the $a_0(980)$ and $a_2(1320)$ region, see Fig. 4, and obtained the following result $\Gamma_{a_0\to\gamma\gamma}=(0.19\pm 0.07^{+0.10}_{-0.07})/B(a_0\to\pi^0\eta)$ keV (about its prehistory see Edwards *at al.*, 1982; Cordier, 1985; Erne, 1985). After, the JADE group (Oest *et al.*, 1990, see also Kolanoski, 1988) obtained $\Gamma_{a_0\to\gamma\gamma}=(0.28{\pm}0.04{\pm}0.10)/B(a_0\to\pi^0\eta)$ keV based on the integral luminosity $149\,\text{pb}^{-1}$ and 291 $\gamma\gamma\to\pi^0\eta$ events. The Crystal Ball and JADE data on the $a_0(980)\to\gamma\gamma$ decay have aroused keen interest (see, for example, Kolanoski, 1986, 1988, 1991; Berger and Wagner, 1987; Kolanoski and Zerwas, 1987; Achasov and Shestakov, 1988; Chanowitz, 1988; Barnes, 1992; Hikasa, *et al.* 1992). Late on, need for high-statistic data arose. But up to now there are no new experiments on the $\gamma\gamma\to\pi^0\eta$ reaction. According to PDG, see Hikasa *et al.* (1992) — Amsler *et al.* (2008), average value for $\Gamma_{a_0\to\gamma\gamma}=(0.24^{+0.8}_{-0.7})/B(a_0\to\pi^0\eta)$ keV.

The JADE group (Oest *et al.*, 1990) measured also the $\gamma\gamma\to\pi^0\pi^0$ cross section and having (60 ± 8)-events in the $f_0(980)$ region[8] obtained for the $f_0\to\gamma\gamma$ decay width $\Gamma_{f_0\to\gamma\gamma}=(0.42\pm 0.06^{+0.08}_{-0.18})$ keV (that corresponds to $\Gamma_{f_0\to\gamma\gamma}<0.6$ keV at 95% C.L.).

Table II First results on the $\gamma\gamma$ width of the $f_0(980)$ (see Field, 1981; Higler, 1981; Edwards *at al.*, 1982; Olsson, 1983; Kolanoski, 1984, 1988).

Experiments	$\Gamma_{f_0\to\gamma\gamma}$ [keV]
TASSO	$(1.3\pm 0.4\pm 0.6)/B(f_0\to\pi^+\pi^-)$
Crystal Ball	$< 0.8/B(f_0\to\pi\pi)$ (95% C.L.)
JADE	< 0.8 (95% C.L.)
Kolanoski, 1988	Average value: 0.27 ± 0.12



Figure 4 Cross section for $\gamma\gamma\to\pi^0\eta$ as function of the invariant mass $\sqrt{s}$ of $\pi^0\eta$. The data are from the Cristal Ball Collaboration (Antreasyan at al., 1986).

In addition, in 1990 the MARK II group in experiment on the $\gamma\gamma\to\pi^+\pi^-$ reaction with the integral luminosity $209\,\text{pb}^{-1}$ (Boyer et al., 1990) and the Crystal Ball group in 1990–1992 in experiments on the $\gamma\gamma\to\pi^0\pi^0$ reaction with the integral luminosities $255\,\text{pb}^{-1}$ (Karch, 1991; Bienlein, 1992) and $97\,\text{pb}^{-1}$ (Marsiske et al., 1990) obtained also similar results for $\Gamma_{f_0\to\gamma\gamma}$. All data are listed together in Table III, and Figs. 5(a) and 5(b) illustrate the manifestations of the $f_0(980)$ and $f_2(1270)$ resonances observed by MARK II and Crystal Ball in the cross sections of the $\gamma\gamma\to\pi\pi$ reactions.

Although the statistical significance of the $f_0(980)$ sig-

Table III 1990–1992 data on the $\gamma\gamma$ width of the $f_0(980)$ (see the text).

Experiments	$\Gamma_{f_0\to\gamma\gamma}$ [keV]
Crystal Ball (1990)	$0.31\pm 0.14\pm 0.09$
MARK II (1990)	$0.29\pm 0.07\pm 0.12$
JADE (1990)	$0.42\pm 0.06^{+0.08}_{-0.18}$
Karch, 1991	0.25 ± 0.10
Bienlein, 1992	$0.20\pm 0.07\pm 0.04$ ≤ 0.31 (90% C.L.)

---

[8] Compare with (2177 ± 47) events which the JADA group had in the $f_2(1270)$ region.



Figure 5 Cross sections for $\gamma\gamma\to\pi^+\pi^-$ (a) and $\gamma\gamma\to\pi^0\pi^0$ (b) as functions of the invariant mass $\sqrt{s}$ of $\pi\pi$. The data correspond to limited angular ranges of the registration of the final pions; θ is the polar angle of the produced π meson in the $\gamma\gamma$ center-of-mass system.

nal in the cross sections and the invariant $\pi\pi$ mass resolution left much to be desired, the existence of a shoulder in the $f_0(980)$ resonance region in the $\gamma\gamma$collision might be thought as established, see Fig.5.

The experiments of eighties and beginning of nineties showed that the two-photon widths of the scalar $f_0(980)$ and $a_0(980)$ resonances are small in comparison with the two-photon widths of the tensor $f_2(1270)$ and $a_2(1320)$ resonances, for which there were obtained the following values $\Gamma_{f_2\to\gamma\gamma}\approx 2.6-3\,\mathrm{keV}$ (Marsiske et al., 1990; Boyer et al., 1990; Oest *et al.*, 1990; Behrend *et al.*, 1992; see also Amsler *et al.*, 2008) and $\Gamma_{a_2\to\gamma\gamma}\approx 1\,\mathrm{keV}$ (Antreasyan *at al.*, 1986; Oest *et al.*, 1990; see also Amsler *et al.*, 2008). This fact pointed to the four-quark nature of the $f_0(980)$ and $a_0(980)$ states (Antreasyan *at al.*, 1986; Kolanoski, 1986; 1988; 1991; Barnes, 1985, 1992; Berger and Wagner, 1987; Kolanoski and Zerwas, 1987; Chanowitz, 1988; Cahn, 1989; Marsiske *et al.*, 1990; Oest *et al.*, 1990; Feindt and Harjes, 1990; Hikasa, *et al.*, 1992).

As mentioned above, in the beginning of eighties it was predicted (Achasov, Devyanin, and Shestakov, 1982a, 1982b) that, if the $a_0(980)$ and $f_0(980)$ mesons are taken as a four-quark states from the lightest nonet of the MIT-bag (Jaffe, 1977), their production rates should be suppressed in photon-photon collisions by a factor ten in relation to the $a_0(980)$ and $f_0(980)$ mesons taken as two-quark P wave states. The estimates obtained for the four-quark model were (Achasov, Devyanin, and Shestakov, 1982a, 1982b)

$$\Gamma_{a_0\to\gamma\gamma}\sim\Gamma_{f_0\to\gamma\gamma}\sim 0.27\,\mathrm{keV}, \tag{3}$$

which were supported by experiments. As for the $q\bar{q}$ model, it predicted that

$$\Gamma_{0^{++}\to\gamma\gamma}/\Gamma_{2^{++}\to\gamma\gamma}=(15/4)\cdot\mathit{corrections}\approx 1.3-5.5 \tag{4}$$

for P wave states with $J^{PC}=0^{++}$ and 2^{++} from the same family (see, for example, Berger and Feld, 1973; Barbieri, Gatto, and Kögerler, 1976; Jackson, 1976; Budnev and Kaloshin, 1979; Barnes, 1985, 1992; Achasov and Shestakov, 1988, Chanowitz, 1988; Marsiske *et al.*, 1990; Morgan and Pennington, 1990; Li, Close, and Barnes, 1991; Kolanoski, 1991; Munz, 1996; Achasov, 1998). This suggested that $\Gamma_{f_0\to\gamma\gamma}\geq 3.4\,\mathrm{keV}$ and $\Gamma_{a_0\to\gamma\gamma}\geq 1.3\,\mathrm{keV}$.

One dwells else on predictions of the molecule model in which the $a_0(980)$ and $f_0(980)$ resonances are nonrelativistic bound states of the $K\bar{K}$ system (Weinstein and Isgur, 1982, 1990). As the $q^2\bar{q}^2$ model, the molecule one explains the state mass degeneracy, their strong coupling with the $K\bar{K}$ channel. As in the four-quark model, in the molecular one no questions arise with the small rates $B[J/\psi\to a_0(980)\rho]/B[J/\psi\to a_2(1320)\rho]$ and $B[J/\psi\to f_0(980)\omega]/B[J/\psi\to f_2(1270)\omega]$ (see specialities in Achasov, 1998, 2002). However, the predictions of this model for the two-photon widths (Barnes, 1985, 1992),

$$\Gamma_{a_0(K\bar{K})\to\gamma\gamma}=\Gamma_{f_0(K\bar{K})\to\gamma\gamma}\approx 0.6\,\mathrm{keV}, \tag{5}$$

are rather big, within two standard deviations contradict the experiment data from Table III. More than that, the widths of $K\bar{K}$ molecules must be smaller (strictly speaking, much smaller) than the binging energy $\epsilon\approx 10$ MeV. Recent data (Amsler *et al.*, 2008), however, contradict this, $\Gamma_{a_0}\sim(50-100)\,\mathrm{MeV}$ and $\Gamma_{f_0}\sim(40-100)\,\mathrm{MeV}$. The $K\bar{K}$ molecule model predicted also (Achasov and Gubin, 1997; Achasov, Gubin, and Shevchenko, 1997) that $B[\phi\to\gamma a_0(980)]\approx B[\phi\to\gamma f_0(980)]\sim 10^{-5}$ that contradicts experiment (Amsler *et al.*, 2008). In addition, recently (Achasov and Kiselev, 2007b, 2008) it was shown that the kaon loop model, ratified by experiment, describes production of a compact state and not an extended molecule. Finally, experiments in which the $a_0(980)$ and $f_0(980)$ mesons were produced in the $\pi^-p\to\pi^0\eta n$ (Dzierba, 1995; Alde *et al.*, 1999) and $\pi^-p\to\pi^0\pi^0 n$ (Alde *et al.*, 1995, 1998; Gunter *et al.*, 2001) reactions within a broad range of four-momentum transfer squares, $0<-t<1\,\mathrm{GeV}^2$, have shown that these states are compact, e.g. as two-quark ρ and other mesons and not as extended molecule ones with form factors determined by the wave functions. These experiments have left no chances for the $K\bar{K}$-molecule model. As to four-quark states, they are as compact as two-quark states [9].

Particle Data Group gives information on an average value of $\Gamma_{f_0\to\gamma\gamma}$ beginning from 1992. Note that no new experimental data on $\Gamma_{f_0\to\gamma\gamma}$ emerged from 1992 up to 2006, nevertheless, its average value, adduced by PDG, evolved noticeably in this period. Based on the data in Table III, the $\Gamma_{f_0\to\gamma\gamma}$ value would be (0.26 ± 0.08) keV. In 1992 PDG obtained the average value $\Gamma_{f_0\to\gamma\gamma}=(0.56\pm0.11)$ keV (Hikasa *et al.*, 1992) combining the JADA (1990) result, see Table III, with the value $\Gamma_{f_0\to\gamma\gamma}=(0.63\pm0.14)$ keV, which was found by Morgan and Pennington (1990) as a result of a theoretical analysis of the MARK II (1990) and Crystal Ball (1990) data. In 1999 Boglione and Pennington carried out a new theoretical analysis of the situation and halved value, $\Gamma_{f_0\to\gamma\gamma}=(0.28\,^{+0.09}_{-0.13})$ keV (see also Pennington, 1999). Particle Data Group noted that the Boglione and Pennington (1999) result replaces the Morgan and Pennington (1990) one but used both results coupled with the JADE (1990) one for calculation of the average $f_0\to\gamma\gamma$ decay width. In this way the value $\Gamma_{f_0\to\gamma\gamma}=(0.39\,^{+0.10}_{-0.13})$ keV emerged in the PDG Review (Groom *et al.*, 2000).

In 2003 a preliminary super-statistics Belle data on $\gamma\gamma\to\pi^+\pi^-$ were reported. They contain the clear signal of the $f_0(980)$ resonance (Mori *et al.*, 2003). In 2005 there emerged our first response (Achasov and Shestakov, 2005) to these data in which the average over the $\pi\pi$ resonance region $<\Gamma_{f_0\to\gamma\gamma}>_{\pi\pi}\approx 0.27$ keV was obtained . It has become clear that $\Gamma_{f_0\to\gamma\gamma}$ is bound to be small. In 2006 PDG excluded the Morgan and Pennington (1990) result, $\Gamma_{f_0\to\gamma\gamma}=(0.63\pm0.14)$ keV, from its sample and using only the JADE (1990) data and the Boglione and Pennington (1999) result obtained a new guide $\Gamma_{f_0\to\gamma\gamma}=(0.31\,^{+0.08}_{-0.11})$ keV (Yao *et al.*, 2006). To the effect that happened later to the average value $\Gamma_{f_0\to\gamma\gamma}$ and can else happens to the one, we are going to tell in the following subsections.

B. Current Experimental Situation

In 2007 the Belle collaboration published the data on cross section of the $\gamma\gamma\to\pi^+\pi^-$ reaction in the region of the $\pi^+\pi^-$ invariant mass, $\sqrt{s}$, from 0.8 up 1.5 GeV based on the integral luminosity $85.9\,\mathrm{fb}^{-1}$ (Mori *et al.*, 2007a, 2007b). These data are shown on Fig. 6. Thanks to the huge statistics and high energy resolution in the Belle experiment, the clear signal of the $f_0(980)$-resonance was detected for the first time. Its value proved to be small that agrees qualitatively with the four-quark model prediction (Achasov and Shestakov, 1982a, 1982b). The visible height of the $f_0(980)$-peak amounts about 15 nb above a smooth background about 100 nb. Its visible (effective) width proved to be about 30–35 MeV, see Fig. 6.



Figure 6 (a) The high statistics Belle data on the $\gamma\gamma\to\pi^+\pi^-$ reaction cross section for $|\cos\theta|\le 0.6$ (Mori *et al.*, 2007b). Plot (b) emphasizes the region of the $f_0(980)$ peak. Errors shown include statistics only. They are approximately equal to 0.5%–1.5%. The $\sqrt{s}$ bin size in the Belle experiment has been chosen to be 5 MeV, with the mass resolution of about 2 MeV.

Table IV The current data on the $\gamma\gamma$ width of the $f_0(980)$.

Experiments	$\Gamma_{f_0\to\gamma\gamma}$ [keV]
Belle (Mori *et al.*, 2007a), $\gamma\gamma\to\pi^+\pi^-$	$0.205^{+0.095+0.147}_{-0.083-0.117}$
PDG (2008) average value	$0.29^{+0.07}_{-0.09}$
Belle (Uehara *et al.*, 2008), $\gamma\gamma\to\pi^0\pi^0$	$0.286\pm0.017^{+0.211}_{-0.070}$

Then the Belle collaboration published the data on cross section for the $\gamma\gamma\to\pi^0\pi^0$ reaction in the region of the $\pi^+\pi^-$ invariant mass, $\sqrt{s}$, from 0.6 to 1.6 GeV based on the integral luminosity $95\,\mathrm{fb}^{-1}$ (Uehara *et al.*, 2008; see also Abe *et al.*, 2007; Nakazawa *et al.*, 2008; Adachi *et al.*, 2008). Here also the clear signal of the $f_0(980)$ resonance was detected for the first time. Note that the background conditions for the manifestation of the $f_0(980)$ in the $\gamma\gamma\to\pi^0\pi^0$ channel are more favourable than in the $\gamma\gamma\to\pi^+\pi^-$ one.

Figures 7(a) and 7(b) illustrate a general picture of data on the cross sections of the $\pi^+\pi^-$ and $\pi^0\pi^0$ production in the photon-photon collisions from the $\pi\pi$ threshold up to 1.5 GeV after the Belle experiments. It is instructive to compare with a previous picture illustrated by Figs. 5(a) and 5(b).

The current information about $\Gamma_{f_0\to\gamma\gamma}$ are adduced in Table IV. The Belle collaboration determined $\Gamma_{f_0\to\gamma\gamma}$ (see Table IV) as a result of fitting the mass distributions (see Figs. 6(b) and 7(b)) taking into account the $f_0(980)$ and $f_2(1270)$ contributions and smooth background contributions, which play the important part in fitting and are a source of large systematic errors in $\Gamma_{f_0\to\gamma\gamma}$ (see details in the papers Mori *et al.*, 2007a and Uehara *et al.*, 2008).

---

[9] A $K\bar{K}$ formation of unknown origin with the relativistic average Euclidean momentum square $<k^2>\approx 2\ \mathrm{GeV}^2$ was considered recently and named "a $K\bar{K}$ molecule" (Branz, Gutsche, Lyubovitskij, 2008). Such a free use of the molecule term can mislead readers considering a molecule as an extent nonrelativistic bound system.



Figure 7 (a) The data on the $\gamma\gamma\to\pi^+\pi^-$ reaction cross section from Mark II (Marsiske *et al.*, 1990) and CELLO (Behrend *et al.*, 1992), for $\sqrt{s}\le 0.85\,$GeV, and from Belle (Mori *et al.*, 2007b), for $0.8\le\sqrt{s}\le 1.5\,$GeV. (b) The data on the $\gamma\gamma\to\pi^0\pi^0$ reaction cross section from Crystal Ball (Boyer *et al.*, 1990), for $\sqrt{s}<0.8\,$GeV, and from Belle (Uehara *et al.*, 2008), for $0.8\le\sqrt{s}\le 1.5\,$GeV. Plots (a), for $\sqrt{s}>0.85\,$GeV, and (b), for $\sqrt{s}>0.8\,$GeV, show exclusively the Belle data to emphasize the discovered miniature signals from the $f_0(980)$ resonance. The theoretical curves, shown on plot (a), correspond to the pure Born cross sections for the process $\gamma\gamma\to\pi^+\pi^-$ for $|\cos\theta|\le0.6$: the total integrated cross section $\sigma^{\rm Born}=\sigma_0^{\rm Born}+\sigma_2^{\rm Born}$ and the integrated cross sections $\sigma_\lambda^{\rm Born}$ with helicity $\lambda=0$ and 2.



Figure 8 The Born diagrams for $\gamma\gamma\to\pi^+\pi^-$.

C. Dynamics of the Reactions $\gamma\gamma\to\pi\pi$

1. Born Contributions and Angular Distributions

To feel the values of the cross sections measured by experiment, in Fig. 7(a) the total Born cross section of the $\gamma\gamma\to\pi^+\pi^-$ process, $\sigma^{\rm Born}=\sigma_0^{\rm Born}+\sigma_2^{\rm Born}$, and the partial helicity ones, $\sigma_\lambda^{\rm Born}$, are adduced as a guide, where $\lambda=0$ or 2 is the absolute value of the photon helicity difference. These cross sections are caused by the point-like one-pion exchange, see Fig. 8.

By the Low theorem[10] and chiral symmetry[11], the Born contributions should dominate near the threshold region of the $\gamma\gamma\to\pi^+\pi^-$-reaction. As shown in Fig. 7(a), this anticipation does not contradict the current data near threshold, but, certainly, errors leave much to be desired. In additional, one can consider the Born contributions as an reasonable approximation of background (non-resonance) contributions in the $\gamma\gamma\to\pi^+\pi^-$ amplitudes in all the resonance region, including the $f_2(1270)$ one. The Born contributions are also the base for a construction of amplitudes, including strong interactions in final state (see, for example, Mennessier, 1983; Lyth, 1984, 1985; Johnson, 1986; Morgan and Pennington, 1987, 1988, 1991; Bijnens and Cornet, 1988; Donoghue, Holstein, and Lin, 1988; Donoghue and Holstein, 1993; Oller and Oset, 1997; Achasov and Shestakov, 2005, 2007; Pennington, 2006).



Figure 9 Plots (a) and (b) show the $\gamma\gamma\to\pi^+\pi^-$ differential cross section in the Born approximation, i.e., in the point-like one-pion exchange approximation, and its components for different values of $\sqrt{s}$/GeV. The vertical straight line $|\cos\theta|=0.6$ shows the upper boundary of the region available for measurements.

The Born contributions have the following particular qualities. First, $\sigma^{\rm Born}$ has a maximum at $\sqrt{s}\approx0.3$ GeV, where $\sigma^{\rm Born}\approx\sigma_0^{\rm Born}$, then $\sigma_0^{\rm Born}$ falls with increasing $\sqrt{s}$, so that the $\sigma_2^{\rm Born}$ contribution dominates in $\sigma^{\rm Born}$ at $\sqrt{s}>0.5$ GeV, see Fig. 7(a). Second, although the $\sigma_2^{\rm Born}$ value is approximately 80 percent caused by the D wave amplitude, its interference with the contribution of higher waves are considerable in the differential cross section $d\sigma^{\rm Born}(\gamma\gamma\to\pi^+\pi^-)/d|\cos\theta|$, compare Figs. 9(a) and 9(b). The interference, destructive in the first half of the $|\cos\theta|\le0.6$ interval and constructive in the second one, flattens out the θ angle distribution in this interval, so that this effect increases with increasing $\sqrt{s}$, see Fig. 9(a).

Since the first resonance with $I^G(J^{PC})=0^+(4^{++})$ has the mass near 2 GeV (Amsler *et al.*, 2008), then seemingly the S and D wave contributions only should dominate at $\sqrt{s}\le1.5$ GeV and the differential cross section of the $\gamma\gamma\to\pi^+\pi^-$ process could be represented as (Mori *et al.*, 2007b)

$$d\sigma(\gamma\gamma\to\pi^+\pi^-)/d\Omega=|S+D_0Y_2^0|^2+|D_2Y_2^2|^2\,, \qquad (6)$$

where S and D_0 (D_2) are the S and D wave amplitudes with the helicity $\lambda=0$ (2), Y_J^m are the spherical harmonics.[12] But, the above discussion shows that the smooth

---

[10] According to this theorem (Low, 1954; Gell-Mann and Goldberger, 1954; Abarbanel and Goldberger, 1968), the Born contributions give the exact physical amplitude of the crossing reaction $\gamma\pi^\pm\to\gamma\pi^\pm$ close to its threshold.

[11] Chiral symmetry guarantees weakness of the $\pi\pi$ interaction at low energy

[12] Eq. (6) corresponds the situation "untagget" when the depen-



Figure 10 The Belle data on the angular distributions for $\gamma\gamma\to\pi^0\pi^0$ (Uehara *et al.*, 2008). The solid lines are the approximations. The vertical straight line $|\cos\theta|=0.8$ shows the upper boundary of the region available for measurements.

background contribution in the $\gamma\gamma\to\pi^+\pi^-$-cross section contains the high partial wave due to the one-pion exchange, so that the smooth background can imitate the large S wave at $|\cos\theta|\leq 0.6$.

The one-pion exchange is absent in the $\gamma\gamma\to\pi^0\pi^0$ channel and the representation of the cross section of this reaction similar to Eq. (6) is a good approximation at $\sqrt{s}\leq 1.5$ GeV

$$d\sigma(\gamma\gamma\to\pi^0\pi^0)/d\Omega=|\widetilde{S}+\widetilde{D}_0Y^0_2|^2+|\widetilde{D}_2Y^2_2|^2\,,\qquad(7)$$

where $\widetilde{S}$ and $\widetilde{D}_0$ ($\widetilde{D}_2$) are the S and D wave amplitudes with the helicity $\lambda=0$ (2). Nevertheless, the partial wave analysis of the $\gamma\gamma\to\pi^0\pi^0$ events, based on Eq. (7), is not prevented from difficulties for the relation $\sqrt{6}|Y^2_2|=Y^0_2-Y^0_0/\sqrt{5}$, which gives no way of separating the partial waves when using only the data on the differential cross section (Oest *et al.*, 1990; Mori *et al.*, 2007b; Uehara *et al.*, 2008). So, the separation of the contributions with the different helicities requests some guesswork, for example, the domination of the helicity 2 in the $f_2(1270)$ resonance production (Krammer and Krasemann, 1978; Krammer, 1978; Krasemann and Vermaseren, 1981; Li, Close, and Barnes, 1991) that agrees rather well with the experimental angle distribution.

The $d\sigma(\gamma\gamma\to\pi^0\pi^0)/d\Omega$ differential cross section in Eq. (7) is a polynomial of the second power of $z=\cos^2\theta$, which can be expressed in terms of its roots z_1 and z_1^*,[13]

$$d\sigma(\gamma\gamma\to\pi^0\pi^0)/d\Omega=C(z-z_1)(z-z_1^*)\,,\qquad(8)$$

where C is a real quantity. So, from fitting experimental data on the differential cross section one can determine only three independent parameters, for example, C, $\mathrm{Re}z_1$ and $\mathrm{Im}z_1$ up to the sign and not four ones, $|\widetilde{S}|$, $|\widetilde{D}_0|$, $|\widetilde{D}_2|$ and $\cos\delta$ (δ is a relative phase between the $\widetilde{S}$ and $\widetilde{D}_0$ amplitudes), as one would like.

In Fig.10 the Belle data on the anglular distributions in $\gamma\gamma\to\pi^0\pi^0$ are adduced at three value of $\sqrt{s}$. All of them are described very well by the simple two-parameter expression $|a|^2+|b\,Y^2_2|^2$ (Achasov and Shestakov, 2008b). This suggests that the $\gamma\gamma\to\pi^0\pi^0$ cross section is saturated only by the $\widetilde{S}$ and $\widetilde{D}_2$ partial wave contributions at $\sqrt{s}<1.5$ GeV.

2. Production Mechanisms of Scalar Resonances

Expectation of the Belle data and their advent have called into being a whole series of theoretical papers which study dynamics of the $f_0(980)$ and $\sigma(600)$ production in the $\gamma\gamma\to\pi\pi$ processes by various means and discuss the nature of these states (Achasov and Shestakov, 2005, 2007, 2008a, 2008b; Pennington, 2006, 2007a, 2007b, 2007c, 2008; Pennington, Mori, Uehara, and Watanabe, 2008; Mennessier, Minkowski, Narison, and Ochs, 2007; Oller, Roca, and Schat, 2008; Oller and Roca, 2008; Mennessier, Narison, and Ochs, 2008a, 2008b; Achasov, Kiselev, and Shestakov, 2008; Achasov, 2008; Mennessier, 2008; Narison, 2008; van Beveren, Kleefeld, and Rupp, 2008; Kalinovsky and Volkov, 2008; Mao, Wang, Zhang, Zheng, and Zhou, 2009).

The main lesson from the analysis of the production mechanisms of the light scalars in the $\gamma\gamma$-collisions is the following (Achasov, 2008a, 2008b). The classical P wave tensor $q\bar{q}$mesons $f_2(1270)$, $a_2(1320)$ and $f'_2(1525)$ are produced in the $\gamma\gamma$ collisions due to the direct $\gamma\gamma\to q\bar{q}$ transitions in the main, whereas the light scalar mesons $\sigma(600)$, $f_0(980)$, and $a_0(980)$ are produced by the rescatterings $\gamma\gamma\to\pi^+\pi^-\to\sigma$, $\gamma\gamma\to K^+K^-\to f_0$, and $\gamma\gamma\to K^+K^-\to a_0$, i.e., due to the four quark transitions. As to the direct transitions $\gamma\gamma\to\sigma$, $\gamma\gamma\to f_0$, and $\gamma\gamma\to a_0$, they are negligible, as it is expected in four-quark model.

This conclusion introduces a new seminal view of the $\gamma\gamma\to\pi\pi$ reaction dynamics at low energy. Let us dwell on this point.

Recall elementary ideas of interactions of C even mesons with photons based on the quark model (Kolanoski, 1984, 1988; Kolanoski and Zerwas, 1987; Amsler *et al.*, 2008). Coupling of the $\gamma\gamma$-system with the classical $q\bar{q}$ states, to which the light pseudoscalar ($J^{PC}=0^{-+}$) and tensor (2^{++}) mesons belong, are proportional to four power of charges of constituent quarks.

Only the width of the $\pi^0\to\gamma\gamma$ decay is evaluated from the first principles (Adler, 1969; Bell and Jackiw, 1969; Bardeen, Fritzsch, and Gell-Mann, 1972; Leutwyler, 1998). $\Gamma_{\pi^0\to\gamma\gamma}$ is determined completely by the Adler-Bell-Jackiw axial anomaly and in this case the theory (QCD) is in excellent agreement with the experiment (Ioffe and Oganesian, 2007; Bernstein, 2007). The relations between the widths of the $\pi^0\to\gamma\gamma$, $\eta\to\gamma\gamma$, and $\eta'\to\gamma\gamma$ decays are obtained in the $q\bar{q}$ model with taking into account the effects of the $\eta-\eta'$ mixing and the $SU(3)$ symmetry breaking (Kolanoski and Zerwas, 1987;

---

dence on the pion azimuth φ is not measured, that took place in all above experiments.

[13] Such a procedure is the base of the determination all solutions when carrying out partial wave analysises, see, for example, Gersten (1969), Barrelet (1972), Petersen (1977), Alde *et al.* (1998, 1999), Sadovsky (1999), Gunter *et al.* (2001).

Leutwyler, 1998; Feldmann, 2000).

As for the tensor mesons, in the ideal mixing case, i.e., if $f_2 = (u\bar{u} + d\bar{d})/\sqrt{2}$ and $f_2' = s\bar{s}$, the quark model predicts the following relations for the coupling constant squares:

$$g^2_{f_2\gamma\gamma} : g^2_{a_2\gamma\gamma} : g^2_{f_2'\gamma\gamma} = 25 : 9 : 2\,. \tag{9}$$

Though absolute values of the two-photon widths of the tensor meson decays cannot be obtained from the first principles (Berger and Feld, 1973; Babcock and Rosner, 1976; Budnev and Kaloshin, 1979; Berger, 1980; Rosner, 1981; Kolanoski, 1984; Berger and Wagner, 1987; Kolanoski and Zerwas, 1987; see also references herein), the $q\bar{q}$ model prediction (9), underlying the relations between the widths of the $f_2(1270)\to\gamma\gamma$, $a_2(1320)\to\gamma\gamma$, and $f_2'(1525)\to\gamma\gamma$ decays, are used with taking into account the effects of a deviation from the ideal mixing and the $SU(3)$ symmetry breaking (Kolanoski, 1984; Poppe, 1986; Kolanoski and Zerwas, 1987; Albrecht *et al.*,1990; Li, Yu, and Shen, 2001; Amsler *et al.*, 2008). Roughly speaking, the $q\bar{q}$ model prediction (9) is borne out by experiment. As for the practically model-independent prediction of the $q\bar{q}$ model $g^2_{f_2\gamma\gamma} : g^2_{a_2\gamma\gamma} = 25 : 9$,[14] this agrees with experiment rather well (Amsler, 2008). This implies that the final state interaction effects are small, in particular, the contributions of the $f_2(1270)\to\pi^+\pi^-\to\gamma\gamma$ rescattering type are small in comparison with the contributions of the direct $q\bar{q}(2^{++})\to\gamma\gamma$ transitions.[15]

The observed smallness of the $a_0(980)$ and $f_0(980)$ meson two-photon widths in comparison with the two-photon tensor meson ones and thus the failure of the $q\bar{q}$ model prediction of the relation (4) between the widths of the direct 0^{++} and $2^{++}\to\gamma\gamma$ transitions point to that $a_0(980)$ and $f_0(980)$ are not the quark and antiquark bound states.[16] If the $q\bar{q}$ component is practically absent in the wave functions of the light scalars and in their $q^2\bar{q}^2$ component the white neutral meson pairs are practically absent too, as in the MIT-bag model, then the $\sigma(600)\to\gamma\gamma$, $f_0(980)\to\gamma\gamma$, and $a_0(980)\to\gamma\gamma$-decays could be the four-quark transitions caused by the rescatterings $\sigma(600)\to\pi^+\pi^-\to\gamma\gamma$, $f_0(980)\to K^+K^-\to\gamma\gamma$ (Achasov, Devyanin, and Shestakov, 1982a, 1982b), and $a_0(980)\to K^+K^-\to\gamma\gamma$. Already in 1998 we considered such a scenario extensively (Achasov and Shestakov, 1988) analyzing the Crystal Ball data (Antreasyan *et al.*, 1986) on the $a_0(980)$ resonance production in the $\gamma\gamma\to\pi^0\eta$ reaction. Fifteen years later, when the preliminary high statics Belle data (Mori *et al.*, 2003) on the $f_0(980)$ resonance production in the $\gamma\gamma\to\pi^+\pi^-$ reaction were reported, we studied what role the rescattering mechanisms, in particular, the $\gamma\gamma\to K^+K^-\to f_0(980)\to\pi^+\pi^-$ mechanism, could play in this process (Achasov and Shestakov, 2005). As a result we showed that just this mechanism gives a reasonable scale of the $f_0(980)$ manifestation in the $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ cross sections.

Then in the $SU(2)_L\times SU(2)_R$ linear σ model frame we showed that the σ field are described by its four-quark component at least in the energy (virtuality) region of the σ resonance and the $\sigma(600)$ meson decay into $\gamma\gamma$ is the four-quark transition $\sigma(600)\to\pi^+\pi^-\to\gamma\gamma$ (Achasov and Shestakov, 2007). We also emphasized that the σ meson contribution in the $\gamma\gamma\to\pi\pi$ amplitudes is shielded due to its strong destructive interference with the background contributions as in the $\pi\pi\to\pi\pi$ amplitudes, i.e., the σ meson is produced in the $\gamma\gamma$ collisions accompanied by the great chiral background due to the rescattering mechanism $\gamma\gamma\to\pi^+\pi^-\to(\sigma + \textit{background})\to\pi\pi$.[17]

The above considerations about dynamics of the $\sigma(600)$, $f_0(980)$, and $f_2(1270)$ resonance production were developed in analyzing the final high-statistics Belle data (Achasov and Shestakov, 2008a, 2008b) on the $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ reactions, to a discussion of which we proceed.

D. Analysis of the New High-Statistics Belle Data

As noted above, the S and $D_{\lambda=2}$ partial wave contributions dominate in the Born cross sections $\sigma_0^{\rm Born}$ and $\sigma_2^{\rm Born}$ respectively in region of interest, $\sqrt{s} < 1.5\,$GeV, and the $\pi\pi$ interaction is strong also in the S and D waves only in this region, that is why the final state strong interaction modifies these Born contribution in $\gamma\gamma\to\pi^+\pi^-$ essentially.[18] In addition, the inelastic $\gamma\gamma\to K^+K^-\to\pi\pi$

---

[14] The deviation from the ideal $f_2(1270)-f_2'(1525)$ mixing increases $g_{f_2\gamma\gamma}$ by less than 3% and the small difference of the $f_2(1270)$ and $a_2(1320)$ masses allows to expect for its small effect also.

[15] Evaluations, see below, show such is indeed the case.

[16] As to the ideal $q\bar{q}$ model prediction $g^2_{f_0\gamma\gamma} : g^2_{a_0\gamma\gamma} = 25 : 9$, this excluded by experiment also.

[17] If it were not for the chiral shielding the $\gamma\gamma\to\pi^0\pi^0$ cross section near the threshold would be $\approx 100\,$nb due to the $\gamma\gamma\to\pi^+\pi^-\to\sigma\to\pi^0\pi^0$ transition (Achasov and Shestakov, 2007) instead of $\approx(5$–$10)$ nb experimentally observed, see Fig. 7(b).

[18] It is reliably established by experiment that the S and D wave contribution dominate in the $\pi\pi$ cross sections in the $I=0$ and 2 channels at $\sqrt{s} < 1.5\,$GeV (see, for example, data Protopopescu *et al.*, 1973; Hyams *et al.*, 1973, 1975; Durusoy *et al.*, 1973; Hoogland *et al.*, 1977; Petersen, 1977; Alde *et al.*, 1995, 1998; Gunter *et al.*, 2001). The partial amplitudes $\pi\pi\to\pi\pi$, $T^I_J(s) = \{\eta^I_J(s)\exp[2i\delta^I_J(s)] - 1\}/[2i\rho_{\pi^+}(s)]$ [where $\delta^I_J(s)$ and $\eta^I_J(s)$ are the phase and inelasticity of the J wave $\pi\pi$ scattering in the channel with the isospin I; $\rho_{\pi^+}(s) = (1-4m^2_{\pi^+}/s)^{1/2}$] with $J=0,2$ and $I=0$ reach their unitarity limits at some value of $\sqrt{s}$ in the region of interest and demonstrate both the smooth energy dependence and the sharp resonance oscillations. The $T^0_2(s)$ amplitude is dominated by the $f_2(1270)$ resonance contribution. The $T^0_0(s)$ amplitude contains the $\sigma_0(600)$ and $f_0(980)$ resonance contributions. The $\sigma_0(600)$ resonance contribution is compensated strongly by the chiral background near the $\pi\pi$



Figure 11 Diagrammatical representation of Eqs. (10) and (11) for amplitudes $\gamma\gamma \to \pi^+\pi^-$.



Figure 12 Diagrammatical representation of Eqs. (12) and (13) for amplitudes $\gamma\gamma \to \pi^0\pi^0$.

rescattering plays the important role in the $f_0(980)$ resonance region (for the first time this process was noted by Achasov, Devyanin, and Shestakov, 1982a, 1982b).

So, we use the model of the helicity, M_λ, and partial, $M_{\lambda J}$, amplitudes of $\gamma\gamma \to \pi\pi$ in which the Born point-like $\pi^\pm$ and $K^\pm$ exchanges modified by the final strong interaction in the S and D_2 waves and the direct transitions of the resonances in two photons are taken into account,[19]

$$\begin{aligned} &M_0(\gamma\gamma \to \pi^+\pi^-; s, \theta) \\ &= M_0^{\text{Born}}(s,\theta) + \widetilde{I}_{\pi^+\pi^-}(s)\, T_{\pi^+\pi^- \to \pi^+\pi^-}(s) \\ &+ \widetilde{I}_{K^+K^-}(s)\, T_{K^+K^- \to \pi^+\pi^-}(s) + M_{\text{res}}^{\text{direct}}(s), \end{aligned} \tag{10}$$

$$\begin{aligned} &M_2(\gamma\gamma \to \pi^+\pi^-; s, \theta) \\ &= M_2^{\text{Born}}(s,\theta) + 80\pi d^2_{20}(\theta) M_{\gamma\gamma \to f_2(1270) \to \pi^+\pi^-}(s), \end{aligned} \tag{11}$$

$$\begin{aligned} &M_0(\gamma\gamma \to \pi^0\pi^0; s, \theta) \\ &= M_{00}(\gamma\gamma \to \pi^0\pi^0; s) = \widetilde{I}_{\pi^+\pi^-}(s)\, T_{\pi^+\pi^- \to \pi^0\pi^0}(s) \\ &+ \widetilde{I}_{K^+K^-}(s)\, T_{K^+K^- \to \pi^0\pi^0}(s) + M_{\text{res}}^{\text{direct}}(s), \end{aligned} \tag{12}$$

$$\begin{aligned} M_2(\gamma\gamma \to \pi^0\pi^0; s, \theta) &= 5 d^2_{20}(\theta) M_{22}(\gamma\gamma \to \pi^0\pi^0; s) \\ &= 80\pi d^2_{20}(\theta) M_{\gamma\gamma \to f_2(1270) \to \pi^0\pi^0}(s) \end{aligned} \tag{13}$$

(here $d^2_{20}(\theta) = (\sqrt{6}/4)\sin^2\theta$).

The diagrams of the above amplitudes are adduced in Figs. 11 and 12.

The first terms in the right sides of Eqs. (10) and (11) are the Born helicity amplitudes $\gamma\gamma \to \pi^+\pi^-$ corresponding the point-like one-pion exchange mechanism (see Fig. 8). Their explicit forms are adduced in Appendix A. The terms in Eqs. (10) and (12), containing the $T_{\pi^+\pi^- \to \pi^+\pi^-}(s) = [2T_0^0(s) + T_0^2(s)]/3$, $T_{\pi^+\pi^- \to \pi^0\pi^0}(s) = 2[T_0^0(s) - T_0^2(s)]/3$, and $T_{K^+K^- \to \pi^+\pi^-}(s) = T_{K^+K^- \to \pi^0\pi^0}(s) = T_{\pi^+\pi^- \to K^+K^-}(s)$ amplitudes, take into account the final state strong interaction in the S wave. Eqs. (10) and (12) imply that $T_{\pi^+\pi^- \to \pi\pi}(s)$ and $T_{K^+K^- \to \pi\pi}(s)$ in the loops, see Figs. 11 and 12, of the $\gamma\gamma \to \pi^+\pi^- \to \pi\pi$ and $\gamma\gamma \to K^+K^- \to \pi\pi$ rescatterings[20] are on the mass shell, resulting in the $\widetilde{I}_{\pi^+\pi^-}(s)$ and $\widetilde{I}_{K^+K^-}(s)$ amplitudes of the triangle diagrams $\gamma\gamma \to \pi^+\pi^- \to$ *a scalar with the mass* $= \sqrt{s}$ and $\gamma\gamma \to K^+K^- \to$ *a scalar with the mass* $= \sqrt{s}$. Their explicit forms are adduced in Appendixes A and C. The amplitude $M_{\text{res}}^{\text{direct}}(s)$ in Eqs. (10) and (12), caused by the direct coupling constants of $\sigma_0(600)$ and $f_0(980)$ with photons, and the $f_2(1270)$ production amplitude $M_{\gamma\gamma \to f_2(1270) \to \pi^+\pi^-}(s) = M_{\gamma\gamma \to f_2(1270) \to \pi^0\pi^0}(s)$ in Eqs. (11) and (13) are specified below.

Let us show by the example of the S-wave amplitudes $M_{00}(\gamma\gamma \to \pi^+\pi^-; s)$ and $M_{00}(\gamma\gamma \to \pi^0\pi^0; s)$ that the unitary condition requirement or the Watson theorem (1952) about interaction in final state holds in the model under consideration. First of all note that the 4π channel contribution is small and consequently $T_{\pi^+\pi^- \to K^+K^-}(s) = e^{i\delta_0^0(s)} |T_{\pi^+\pi^- \to K^+K^-}(s)|$ and $M_{\text{res}}^{\text{direct}}(s) = \pm e^{i\delta_0^0(s)} |M_{\text{res}}^{\text{direct}}(s)|$ at $4m_\pi^2 \le s \le 4m_K^2$ (Achasov and Shestakov, 2005, 2007, 2008a; Achasov and Kiselev, 2006, 2007a). Taking into account that

---

threshold to provide for the observed smallness of the $\pi\pi$ scattering length a_0^0 and the Adler zero in $T_0^0(s)$ at $s \approx m_\pi^2/2$ (Achasov and Shestakov, 1994a, 2007; Achasov and Kiselev, 2006, 2007a). $|T_0^0(s)|$ reaches the unitary limit in the 0.85–0.9 GeV region and has the narrow deep (practically up to zero) right under the $K\bar{K}$ threshold caused by the destructive interference of the $f_0(980)$ resonance contribution with the large smooth background. It is established also that the $\pi\pi$ scattering in the $I = 0$ channel is elastic up to the $K\bar{K}$ channel threshold in the very good approximation, but directly above this the inelasticity $\eta_0^0(s)$ shows the sharp jump due to the production of the $f_0(980)$ resonance coupled strongly with the $K\bar{K}$channel.

[19] See, in addition, Mennessier, 1983; Achasov and Shestakov, 1988, 1992, 1994b, 2005, 2007, 2008a, 2008b; Achasov and Gubin, 1998; Pennington, Mori, Uehara, and Watanabe, 2008; Oller and Roca, 2008; Mennessier, Narison, and Ochs, 2008a.

[20] The Born diagrams for $\gamma\gamma \to K\bar{K}$ are the same ones which are in Fig. 8 but with replacing the pions by the kaons. The corresponding helicity amplitudes are in Appendix C.

$\mathrm{Im}\widetilde{I}_{\pi^+\pi^-}(s)=\rho_{\pi^+}(s)M_{00}^{\mathrm{Born}}(s)$ one finds

$$\begin{aligned} M_{00}(\gamma\gamma\to\pi^+\pi^-;s) &= M_{00}^{\mathrm{Born}}(s) \\ &+\widetilde{I}_{\pi^+\pi^-}(s)T_{\pi^+\pi^-\to\pi^+\pi^-}(s) \\ &+\widetilde{I}_{K^+K^-}(s)T_{K^+K^-\to\pi^+\pi^-}(s)+M_{\mathrm{res}}^{\mathrm{direct}}(s) \\ &= (\text{for } 2m_\pi\le\sqrt{s}\le 2m_K) \\ &= \tfrac{2}{3}e^{i\delta_0^0(s)}A(s)+\tfrac{1}{3}e^{i\delta_0^2(s)}B(s), \end{aligned} \tag{14}$$

$$\begin{aligned} M_{00}(\gamma\gamma\to\pi^0\pi^0;s) &= \widetilde{I}_{\pi^+\pi^-}(s)T_{\pi^+\pi^-\to\pi^0\pi^0}(s) \\ &+\widetilde{I}_{K^+K^-}(s)T_{K^+K^-\to\pi^0\pi^0}(s)+M_{\mathrm{res}}^{\mathrm{direct}}(s) \\ &= (\text{for } 2m_\pi\le\sqrt{s}\le 2m_K) \\ &= \tfrac{2}{3}e^{i\delta_0^0(s)}A(s)-\tfrac{2}{3}e^{i\delta_0^2(s)}B(s), \end{aligned} \tag{15}$$

where A(s) and B(s) the real functions.[21] Eqs. (14) and (15) show that at one with the Watson theorem the phases of the S wave amplitudes $\gamma\gamma\to\pi\pi$ with $I=0$ and 2 coincide with the phases of the $\pi\pi$ scattering $\delta_0^0(s)$ and $\delta_0^2(s)$, respectively, in the elastic region (below the $K\bar{K}$ threshold).

We use the following notation and normalization

$$\sigma(\gamma\gamma\to\pi^+\pi^+;|\cos\theta|\le 0.6)\equiv\sigma=\sigma_0+\sigma_2, \tag{16}$$

$$\sigma(\gamma\gamma\to\pi^0\pi^0;|\cos\theta|\le 0.8)\equiv\tilde{\sigma}=\tilde{\sigma}_0+\tilde{\sigma}_2, \tag{17}$$

$$\sigma_\lambda=\tfrac{\rho_{\pi^+}(s)}{64\pi s}\int_{-0.6}^{0.6}|M_\lambda(\gamma\gamma\to\pi^+\pi^-;s,\theta)|^2 d\cos\theta, \tag{18}$$

$$\tilde{\sigma}_\lambda=\tfrac{\rho_{\pi^+}(s)}{128\pi s}\int_{-0.8}^{0.8}|M_\lambda(\gamma\gamma\to\pi^0\pi^0;s,\theta)|^2 d\cos\theta. \tag{19}$$

$\sigma_{\lambda J}$ and $\tilde{\sigma}_{\lambda J}$ denote the corresponding partial cross sections.

Before fitting data it is helpful to center on a simplified (qualitative) scheme of their description.

In Fig. 13(a) (from Achasov and Shestakov, 2008a) are adduced the theoretical curves for the cross section $\sigma=\sigma_0+\sigma_2^{\mathrm{Born}}$ and its components σ_0 and σ_2^{Born} corresponding to the simplest variant of the above model in which only the S wave Born amplitudes $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to K^+K^-$ are modified by the pion and kaon final state strong interaction[22]. This modification results in appearing the $f_0(980)$ resonance signal in σ_0, the value and shape of which agree very well with the Belle data, see Fig. 13(a). From comparing the corresponding curves in Figs. 13(a) and 7(a) it follows that the S wave contribution to $\sigma(\gamma\gamma\to\pi^+\pi^-;|\cos\theta|\le 0.6)$ is small at $\sqrt{s}>0.5$ GeV in any case. It is clear that the $f_2(1270)$ resonance contribution is the main element required for the description of the Belle data on $\gamma\gamma\to\pi^+\pi^-$ in the $\sqrt{s}$ region from 0.8 up to 1.5 GeV. For describing data only near the $f_0(980)$ resonance one can the large non-coherent background under the resonance, caused by σ_2, approximate by a polynomial of $\sqrt{s}$. The result of a such fit is shown in Figs. 13(c) and 13(d) (Achasov and Shestakov, 2008a).



Figure 13 Theoretical curves in plots (a) and (b) correspond to the simplest model which incorporates only the Born contributions $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to K^+K^-$ modified for strong final-state interactions in the S wave. Plot (c) illustrates the description of the Belle data in the $f_0(980)$ region (see the text for details); (d) is a fragment of (c).

By Fig. 12 and Eq. (12) taking into account the final state interaction in the Born $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to K^+K^-$ amplitudes leads to the prediction of the S-wave amplitude of the $\gamma\gamma\to\pi^0\pi^0$ reaction (Achasov and Shestakov, 2005, 2007, 2008a). In Figs. 13(b) the $\gamma\gamma\to\pi^0\pi^0$ cross section, $\tilde{\sigma}_0=\tilde{\sigma}_{00}$, evaluated in the outlined above manner, are compared with the Crystal Ball and Belle data.[23] In view of the fact that no fitting parameters are used for the construction of $\tilde{\sigma}_0$, one should accept that the agreement with the data is rather well at $\sqrt{s}\le 0.8$ Gev, i.e., in the $\sigma(600)$ resonance region. It is clear also that at $\sqrt{s}>0.8$ GeV the $f_2(1270)$ resonance responsibility region begins.

So, already at this stage it emerges the following. First, if the direct coupling constants of $\sigma(600)$ and $f_0(980)$ with $\gamma\gamma$ are included in fitting their role will

---

[21] $A(s)=M_{00}^{\mathrm{Born}}(s)\cos\delta_0^0(s)+(1/\rho_{\pi^+}(s))\mathrm{Re}[I_{\pi^+\pi^-}(s)]\sin\delta_0^0(s)+(3/2)\widetilde{I}_{K^+K^-}(s)|T_{K^+K^-\to\pi^+\pi^-}(s)|\pm|M_{\mathrm{res}}^{\mathrm{direct}}(s)|$ and $B(s)=M_{00}^{\mathrm{Born}}(s)\cos\delta_0^2(s)+(1/\rho_{\pi^+}(s))\mathrm{Re}[\widetilde{I}_{\pi^+\pi^-}(s)]\sin\delta_0^2(s)$.

[22] As for all higher partial waves with $\lambda=0$ and 2, they are taken in the Born point-like approximation (Achasov and Shestakov, 2005, 2008a).

[23] Note that the step of $\sqrt{s}$ for the Crystal Ball and Belle data, shown in Fig. 13(b), is 50 MeV and 20 MeV respectively.



Figure 14 The structure of the $f_0(980)$ signal in σ_0. (a) The contributions from the $\gamma\gamma \to K^+K^- \to \pi^+\pi^-$ (dashed line), $\gamma\gamma \to \pi^+\pi^- \to \pi^+\pi^-$ (dotted line) rescattering amplitudes, and their sum (solid line). (b) The dashed line is identical to the solid one in (a), the dotted and dot-dashed lines show the $\sigma_0^{\rm Born}$ and $\sigma_{00}^{\rm Born}$ cross sections, respectively ($\sigma_0^{\rm Born} < \sigma_{00}^{\rm Born}$ because of the destructive interference between the S and higher partial waves), and the solid line corresponds to the resulting $f_0(980)$ signal in σ_0.

be negligible in agreement with the four-quark model prediction (Achasov, Devyanin, and Shestakov, 1982a, 1982b). Second, by Eqs. (10) and (12) the $\sigma(600)\to\gamma\gamma$ and $f_0(980)\to\gamma\gamma$ decays are described by the triangle loop rescattering diagrams $Resonance \to (\pi^+\pi^-, K^+K^-) \to \gamma\gamma$ and, consequently, are the four-quark transitions (Achasov and Shestakov, 2005, 2007, 2008a, 2008b).

The interesting and important feature of the $f_0(980)$ signal in $\gamma\gamma\to\pi^+\pi^-$ is its complicated structure which is shown by Figs. 14(a) and 14(b). The $\gamma\gamma\to K^+K^-\to\pi\pi$ rescattering amplitude plays the determinant role transferring the $f_0(980)$ peak from the $T_{K^+K^-\to\pi\pi}(s)$ amplitude to the $\gamma\gamma\to\pi\pi$ one. The $\gamma\gamma\to\pi^+\pi^-\to\pi^+\pi^-$ rescattering in its turn transfers the narrow deep under the $K\bar{K}$ threshold from the $T_{\pi\pi\to\pi\pi}(s)$ amplitude, see the footnote 18, to the $\gamma\gamma\to\pi\pi$ one. The interference of the resonance $\gamma\gamma\to K^+K^-\to\pi^+\pi^-$ amplitude with the $\gamma\gamma\to\pi^+\pi^-\to\pi^+\pi^-$ amplitude [24] exerts the essential effect on the resulting shape of the $f_0(980)$ signal, as indicated by Fig. 14(a). As for the Born contributions, their influence on on the resulting shape of the $f_0(980)$ signal is small, see Fig. 14(b).

Once more notable fact lies in the drastic change of the $f_0(980)$ production amplitude $\gamma\gamma\to K^+K^-\to f_0(980)$ in the $f_0(980)$ peak region (Achasov and Shestakov, 2005, 2008a, 2008b) just as the $\gamma\gamma\to K^+K^-\to a_0(980)$ amplitude in the $a_0(980)$ region (Achasov and Shestakov, 1988), see Section IV. A. In the cross section its contribution is proportional to $|\widetilde{I}_{K^+K^-}(s)|^2$, see Eqs. (10) and (12). The function $|\widetilde{I}_{K^+K^-}(s)|^2$ decreases drastically immediately under the K^+K^- threshold,i.e., in the $f_0(980)$ resonance region, see Fig. 15.[25] Such a behavior of the $f_0(980)$ two-photon production amplitude reduces strongly the left slope of the the $f_0(980)$ peak defined by the resonance amplitude $T_{K^+K^-\to\pi\pi}(s)$. That is why one cannot approximate the $f_0(980)\to\gamma\gamma$ decay width by a constant even in the region $m_{f_0} - \Gamma_{f_0}/2 \le \sqrt{s} \le m_{f_0} + \Gamma_{f_0}/2$ (Achasov and Shestakov, 2005).



Figure 15 The solid curve shows $|\widetilde{I}_{K^+K^-}(s)|^2$ as a function of $\sqrt{s}$ (see Appendix. A); the dashed and dotted curves above the K^+K^- threshold correspond to the contributions of the real and imaginary parts of $\widetilde{I}_{K^+K^-}(s)$, respectively.

So, the above consideration teaches us that all simplest approximations of the $f_0(980)$ signal shape observed in the $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ cross sections can give only a rather relative information on the $f_0(980)$ state two-photon production mechanism and the $f_0(980)$ parameters.

Fortunately, the already current knowledge of the dynamics of $T_{\pi^+\pi^-\to\pi\pi}(s)$ $[T_0^0(s), T_0^2(s)]$ and $T_{K^+K^-\to\pi\pi}(s)$ strong interaction amplitudes allows to advance in understanding the signals about the light scalar mesons which the data on the $\gamma\gamma\to\pi\pi$ reaction send to us. In fitting data we use the model for the $T_0^0(s)$ and $T_{K^+K^-\to\pi\pi}(s)$ amplitudes which was suggested and used for the joint analysis of the data on the $\pi^0\pi^0$ spectrum mass in the $\phi\to\pi^0\pi^0\gamma$ decay, the $\pi\pi$ scattering at $2m_\pi < \sqrt{s} < 1.6$ GeV and the $\pi\pi\to K\bar{K}$ reaction (Achasov and Kiselev, 2006, 2007a). The $T_0^0(s)$ model takes into account the contributions of the $\sigma(600)$ and $f_0(980)$ resonances, their mixing, and the chiral background with the large negative phase which shields the $\sigma(600)$ resonance (see additionally Achasov and Shestakov, 1994a, 2007). Eqs. (10) and (12) transfer the effect of the chiral shielding of the $\sigma(600)$ resonance from the $\pi\pi$ scattering into the $\gamma\gamma\to\pi\pi$ amplitudes. This effect are demonstrated by Fig. 16(a) with the help of the $\pi\pi$ scattering phases $\delta_{\rm res}(s)$, $\delta_B^{\pi\pi}(s)$, and $\delta_0^0(s)$ [see Eqs. (20) − (22) and (24)], and by Figs. 16(b) and 16(c) with the help the corresponding cross sections of the $\pi\pi\to\pi\pi$ and $\gamma\gamma\to\pi^0\pi^0$ reactions (Achasov and Shestakov, 2007). As seen from Fig.16(c), if it were not for such a shielding the $\gamma\gamma\to\pi^0\pi^0$ cross section nearby the threshold would be not $5-10$ nb but approximately 100 nb due to the $\pi^+\pi^-$ loop mechanism of the decay $\sigma(600)\to\gamma\gamma$ (Achasov and Shestakov, 2007). The width decay corresponding to this mechanism, $\Gamma_{\sigma\to\pi^+\pi^-\to\gamma\gamma}(s)$, is shown in Fig. 16(d), see also Eq. (51) in Appendix A.

---

[24] Note that the relative sign between these amplitudes is fixed surely. (Achasov and Shestakov, 2005, 2008a, 200b).

[25] The function $|\widetilde{I}_{K^+K^-}(s)|^2$ decreases relatively its maximum at $\sqrt{s}=2m_{K^+}\approx 0.9873$ GeV by 1.66, 2.23, 2.75, 3.27, and 6.33 times at $\sqrt{s}=0.98$, 0.97, 0.96, 0.95, and 0.9 GeV, respectively.

By Achasov and Kiselev (2006, 2007b),

$$T_0^0(s)=T_B^{\pi\pi}(s)+e^{2i\delta_B^{\pi\pi}(s)}T_{\rm res}^{\pi\pi}(s)\,, \qquad (20)$$

$$T_B^{\pi\pi}(s)=\{\exp[2i\delta_B^{\pi\pi}(s)]-1\}/[2i\rho_{\pi^+}(s)]\,, \qquad (21)$$

$$T_{\rm res}^{\pi\pi}(s)=\{\eta_0^0(s)\exp[2i\delta_{\rm res}(s)]-1\}/[2i\rho_{\pi^+}(s)]\,, \qquad (22)$$

$$T_{K^+K^-\to\pi^+\pi^-}(s)=e^{i[\delta_B^{\pi\pi}(s)+\delta_B^{K\bar K}(s)]}T_{\rm res}^{K\bar K\to\pi\pi}(s)\,, \qquad (23)$$

where $\delta_B^{\pi\pi}(s)$ and $\delta_B^{K\bar K}(s)$ are the phase of the elastic S wave background in the $\pi\pi$ and $K\bar K$ channels with $I=0$;

$$\delta_0^0(s)=\delta_B^{\pi\pi}(s)+\delta_{\rm res}(s). \qquad (24)$$

The amplitudes of the $\sigma(600)-f_0(980)$ resonance complex in Eqs. (10), (12), (20), (22), and (23) are (Achasov and Kiselev, 2006, 2007b)

$$T_{\rm res}^{\pi\pi}(s)=3\,\frac{g_{\sigma\pi^+\pi^-}\Delta_{f_0}(s)+g_{f_0\pi^+\pi^-}\Delta_\sigma(s)}{32\pi[D_\sigma(s)D_{f_0}(s)-\Pi^2_{f_0\sigma}(s)]}\,, \qquad (25)$$

$$T_{\rm res}^{K\bar K\to\pi\pi}(s)=\frac{g_{\sigma K^+K^-}\Delta_{f_0}(s)+g_{f_0K^+K^-}\Delta_\sigma(s)}{16\pi[D_\sigma(s)D_{f_0}(s)-\Pi^2_{f_0\sigma}(s)]}\,, \qquad (26)$$

$$M_{\rm res}^{\rm direct}(s)=s\,e^{i\delta_B^{\pi\pi}(s)}\,\frac{g^{(0)}_{\sigma\gamma\gamma}\Delta_{f_0}(s)+g^{(0)}_{f_0\gamma\gamma}\Delta_\sigma(s)}{D_\sigma(s)D_{f_0}(s)-\Pi^2_{f_0\sigma}(s)}\,, \qquad (27)$$

where $\Delta_{f_0}(s)=D_{f_0}(s)g_{\sigma\pi^+\pi^-}+\Pi_{f_0\sigma}(s)g_{f_0\pi^+\pi^-}$ and $\Delta_\sigma(s)=D_\sigma(s)g_{f_0\pi^+\pi^-}+\Pi_{f_0\sigma}(s)g_{\sigma\pi^+\pi^-}$; $g^{(0)}_{\sigma\gamma\gamma}$ and $g^{(0)}_{f_0\gamma\gamma}$ are the direct coupling constants of the σ and f_0 resonances with the photons. We use the expressions for the $\delta_B^{\pi\pi}(s)$ and $\delta_B^{K\bar K}(s)$ phases, the propagators $\sigma(600)$ and $f_0(980)$ resonances $1/D_\sigma(s)$ and $1/D_{f_0}(s)$ and the polarization operator matrix element $\Pi_{f_0\sigma}(s)$ from (Achasov and Kiselev, 2006). The m_{f_0} was free, the other parameters in the strong amplitudes (m_σ, $g_{\sigma\pi^+\pi^-}$, $g_{f_0K^+K^-}$, etc.) correspond to the variant 1 from Table 1 from this paper.[26] We also put $\eta_0^2(s)=1$ for all $\sqrt{s}$ under consideration an take $\delta_0^2(s)$ from (Achasov and Shestakov, 2003).



Figure 16 The figure demonstrates the chiral shielding effect in the reactions $\pi\pi\to\pi\pi$ and $\gamma\gamma\to\pi^0\pi^0$. All the plots have been taken from the paper Achasov and Shestakov (2007) dedicated the lightest scalar in the $SU(2)_L\times SU(2)_R$ linear σ model.

The amplitudes of the $f_2(1270)$ resonance production in Eqs. (11) and (13) are

$$M_{\gamma\gamma\to f_2(1270)\to\pi^+\pi^-}(s)=M_{\gamma\gamma\to f_2(1270)\to\pi^0\pi^0}(s)$$

$$=\frac{\sqrt{s}\,G_2(s)\sqrt{2\Gamma_{f_2\to\pi\pi}(s)/3}}{m^2_{f_2}-s-i\sqrt{s}\Gamma^{\rm tot}_{f_2}(s)}\,. \qquad (28)$$

The main contribution in its total width $\Gamma^{\rm tot}_{f_2}(s)=\Gamma_{f_2\to\pi\pi}(s)+\Gamma_{f_2\to K\bar K}(s)+\Gamma_{f_2\to4\pi}(s)$ is given by the $\pi\pi$ partial decay width

$$\Gamma_{f_2\to\pi\pi}(s)=\Gamma^{\rm tot}_{f_2}(m^2_{f_2})B(f_2\to\pi\pi)$$

$$\times\frac{m^2_{f_2}}{s}\,\frac{q^5_{\pi^+}(s)}{q^5_{\pi^+}(m^2_{f_2})}\,\frac{D_2(q_{\pi^+}(m^2_{f_2})R_{f_2})}{D_2(q_{\pi^+}(s)R_{f_2})}\,, \qquad (29)$$

where $D_2(x)=9+3x^2+x^4$, $q_{\pi^+}(s)=\sqrt{s}\rho_{\pi^+}(s)/2$, R_{f_2} is the interaction range and $B(f_2\to\pi\pi)=0.848$ (Amsler *et al.*, 2008). The small contributions of $\Gamma_{f_2\to K\bar K}(s)$ and $\Gamma_{f_2\to4\pi}(s)$ are the same ones as in (Achasov and Shestakov, 2008a). The parameter R_{f_2} (Marsiske *et al.*, 1990; Boyer *et al.*, 1990; Behrend *et al.*, 1992; Mori *et al.*, 2007b; Uehara *et al.*, 2008; Achasov and Shestakov, 2008a, 2008b) controls the relative form of the $f_2(1270)$-resonance wings and is very important especially for fitting data with small errors.

---

[26] Removing the misprint in the sign of the constant $C\equiv C_{f_0\sigma}$ we use $C_{f_0\sigma}=-0.047$ GeV. Notice that our principal conclusions [the insignificance of the direct transition $\gamma\gamma\to$ *Light Scalar* and the dominant role of the four-quark transition $\gamma\gamma\to(\pi\pi,\,K^+K^-)\to$ *Light Scalar*] are independent on a specific variant from (Achasov and Kiselev, 2006, 2007b).

The amplitude $G_2(s)$ in Eq. (28) describe the coupling of the $f_2(1270)$-resonance with the photons,

$$G_2(s) = \sqrt{\Gamma^{(0)}_{f_2\to\gamma\gamma}(s)} + i\frac{M^{\rm Born}_{22}(s)}{16\pi}\sqrt{\frac{2}{3}\rho_{\pi^+}(s)\Gamma_{f_2\to\pi\pi}(s)}\,. \tag{30}$$

The explicit form of the $M^{\rm Born}_{22}(s)$ amplitude is in Appendix A, Eq. (41). The $f_2(1270)\to\gamma\gamma$ decay width is

$$\Gamma_{f_2\to\gamma\gamma}(s) = |G_2(s)|^2 \tag{31}$$

and

$$\Gamma^{(0)}_{f_2\to\gamma\gamma}(s) = \frac{m_{f_2}}{\sqrt{s}}\Gamma^{(0)}_{f_2\to\gamma\gamma}(m^2_{f_2})\frac{s^2}{m^4_{f_2}} \tag{32}$$

[here the s^2 factor, and also the s factor in Eq. (27), is caused by the gauge invariance requirement]. The second term in $G_2(s)$ corresponds to the rescattering $f_2(1270)\to\pi^+\pi^-\to\gamma\gamma$ with the real pions in intermediate state[27] and ensure the fulfillment of the Watson theorem requirement for the amplitude $\gamma\gamma\to\pi\pi$ with $\lambda = J = 2$ and $I = 0$ under the first inelastic threshold. This term gives a small contribution, less then 6%, in $\Gamma_{f_2\to\gamma\gamma}(m^2_{f_2})$.[28]

The simplest approximation (32) of the main contribution, $\Gamma^{(0)}_{f_2\to\gamma\gamma}(s)$, in the $f_2(1270)\to\gamma\gamma$ decay width is completely adequate to the current state of both theory and experiment. The parameter $\Gamma^{(0)}_{f_2\to\gamma\gamma}(m^2_{f_2}) = \frac{1}{5}[g^2_{f_2\gamma\gamma}/(16\pi)]m^3_{f_2}$ in Eq. (32) accumulates effectively lack of knowledge of the values of the amplitudes responsible for the $f_2(1270)\to\gamma\gamma$ decay. By the above reasons, see Subsection III. C. 2., it is generally agreed that the direct quark-antiquark transition $q\bar{q}\to\gamma\gamma$ dominates in the $f_2(1270)\to\gamma\gamma$ decay and its amplitude is characterized by the $g_{f_2\gamma\gamma}$ coupling constant. As shown in (Achasov and Shestakov, 1988, 2005, 2007, 2008a, 2008b) and as we state here step by step, the situation is quite different in the case of the light scalar mesons.

Now everything is ready to come to the discussion of fitting the Belle data on the $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ cross sections which was carried out in (Achasov and Shestakov, 2008a, 2008b).

One considers firstly fitting the $\gamma\gamma\to\pi^0\pi^0$ cross section only, see Fig. 17(b), which has the smaller background contributions under the $f_0(980)$ and $f_2(1270)$ resonances then the $\gamma\gamma\to\pi^+\pi^-$ cross section, compare Figs. 17(a) and 17(b). The solid curve in Fig. 17(b), describing these data rather well, corresponds the following parameters of the model: $m_{f_2} = 1.269$ GeV, $\Gamma^{\rm tot}_{f_2}(m^2_{f_2}) = 0.182$ GeV, $R_{f_2} = 8.2$ GeV^{-1}, $\Gamma_{f_2\to\gamma\gamma}(m_{f_2}) = 3.62$ keV $[\Gamma^{(0)}_{f_2\to\gamma\gamma}(m_{f_2}) = 3.43$ keV], $m_{f_0} = 0.969$ GeV, $g^{(0)}_{\sigma\gamma\gamma} = 0.536$ GeV^{-1} and $g^{(0)}_{f_0\gamma\gamma} = 0.652$ GeV^{-1}.[29] The fitting indicates smallness of the direct coupling constants $g^{(0)}_{\sigma\gamma\gamma}$ and $g^{(0)}_{f_0\gamma\gamma}$: $\Gamma^{(0)}_{\sigma\to\gamma\gamma}(m^2_\sigma) = |m^2_\sigma g^{(0)}_{\sigma\gamma\gamma}|^2/(16\pi m_\sigma) = 0.012$ keV and $\Gamma^{(0)}_{f_0\to\gamma\gamma}(m^2_{f_0}) = |m^2_{f_0} g^{(0)}_{f_0\gamma\gamma}|^2/(16\pi m_{f_0}) = 0.008$ keV, in accordance with the prediction (Achasov, Devyanin, and Shestakov, 1982a, 1982b).[30] The dominant rescattering mechanisms give the $\sigma(600)\to\pi^+\pi^-\to\gamma\gamma$ width $\approx(1-1.75)$ keV in the region $0.4 < \sqrt{s} < 0.5$ GeV, see Fig. 16(d) (Achasov and Shestakov, 2007), and the $f_0(980)\to K^+K^-\to\gamma\gamma$ width $\approx(0.15-0.2)$ keV averaged over the resonance mass distibution (Achasov and Shestakov, 2005).



Figure 17 Cross sections for the $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ reactions. Only statistical errors are shown for the Belle data (Mori *et al.*, 2007b; Uehara *et al.*, 2008). The curves in plot (a) are described in the text and on the figure. The curves in plot (b) are the result of the fit to the data on the $\gamma\gamma\to\pi^0\pi^0$ reaction.

But such a fitting of the $\gamma\gamma\to\pi^0\pi^0$ cross section comes into conflict with the data on $\gamma\gamma\to\pi^+\pi^-$, see solid curve for $\sigma = \sigma_0 + \sigma_2$ in Fig. 17(a). This is connected with the large Born contribution in σ_2 and its strong constructive (destructive) interference with the $f_2(1270)$ resonance contribution at $\sqrt{s} < m_{f_2}$ ($\sqrt{s} > m_{f_2}$), which are absent in $\gamma\gamma\to\pi^0\pi^0$. We faced this challenge in (Achasov and Shestakov, 2008a) and ibidem suggested the following solution. Matters can be improved by the introduction of the common cutting form factor in the

---

[27] That is, it corresponds to the imaginary part of the $f_2(1270)\to\pi^+\pi^-\to\gamma\gamma$ amplitude.

[28] As for a real part of the $f_2(1270)\to\pi^+\pi^-\to\gamma\gamma$ amplitude, its modulus is far less than the one of the direct transition amplitude as different estimations show.

[29] The formally calculated errors in the significant parameters of the model are negligible due to the high statistical accuracy of the Belle data. The model dependence of adjustable parameter values is the main source of their ambiguity.

[30] One notes that the small values of these coupling constants are grasped in fitting due to the interference of the $M^{\rm direct}_{\rm res}(s)$ amplitude, see Eqs. (10), (12), and (27), with the contributions of the dominant rescattering mechanisms. In such a case not the specific above values of $g^{(0)}_{\sigma\gamma\gamma}$ and $g^{(0)}_{f_0\gamma\gamma}$ are important, but the fact of their relative smallness corresponding $\Gamma^{(0)}_{\sigma\to\gamma\gamma}(m^2_\sigma)$ and $\Gamma^{(0)}_{f_0\to\gamma\gamma}(m^2_{f_0}) \ll 0.1\,$keV.



Figure 18 Joint description of the data on the cross sections for the reactions $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$. The shaded bands correspond to the Belle data (Mori *et al.*, 2007b; Uehara *et al.*, 2008) with the statistical and systematic errors (errors are added quadratically). The curves are described in the text and on the figures; ${\sigma_2}^{\rm Born}(x_1)$ in plot (a) is the Born cross section for the $\gamma\gamma\to\pi^+\pi^-$ reaction with the inclusion of the form factor.

point-like Born amplitudes $\gamma\gamma\to\pi^+\pi^-$, $M^{\rm Born}_\lambda(s,\theta)\to G(t,u)M^{\rm Born}_\lambda(s,\theta)$, where t and u are the Mandelstam variables for the $\gamma\gamma\to\pi^+\pi^-$ reaction.[31] To show this we use, as an example, the expression for $G(t,u)$ suggested by Poppe (1986)

$$G(t,u)=\frac{1}{s}\left[\frac{m^2_{\pi^+}-t}{1-(u-m^2_{\pi^+})/x_1^2}+\frac{m^2_{\pi^+}-u}{1-(t-m^2_{\pi^+})/x_1^2}\right], \tag{33}$$

where x_1 is a free parameter. This ansatz is quite acceptable in the physical region of the $\gamma\gamma\to\pi^+\pi^-$reaction. Replacing m_{π^+} by m_{K^+} and x_1 by x_2 we obtain also the form factor for the Born amplitudes $\gamma\gamma\to K^+K^-$.

The solid curves for $\sigma=\sigma_0+\sigma_2$ and $\tilde\sigma=\tilde\sigma_0+\tilde\sigma_2$ in Fig. 18(a) and 18(b) show the consistent fitting of the data on the $\gamma\gamma\to\pi^+\pi^-$ cross section in the region $0.85<\sqrt{s}<1.5\,$GeV and on the $\gamma\gamma\to\pi^0\pi^0$ cross section in the region $2m_\pi<\sqrt{s}<1.5\,$GeV with the form factors modificating the point-like Born contributions. The obtained description is more than satisfactory to within the Belle systematic errors which are shown in Figs. 18(a) and 18(b) by means of the shaded bands. We believe that such a fitting is completely adequate for the statistic errors of both Belle measurements are so small that to obtain the formally good enough value of χ^2 in the combined fitting of the $\pi^+\pi^-$ and $\pi^0\pi^0$ data in the wide regions of $\sqrt{s}$ without taking the systematic errors into consideration is practically impossible.[32] The curves in Fig. 18 corresponds the following values of the parameters: $m_{f_2}=1.272\,$GeV, $\Gamma^{\rm tot}_{f_2}(m^2_{f_2})=0.196\,$GeV, $R_{f_2}=8.2\,{\rm GeV}^{-1}$, $\Gamma_{f_2\to\gamma\gamma}(m_{f_2})=3.83\,$keV $[\Gamma^{(0)}_{f_2\to\gamma\gamma}(m_{f_2})=3.76\,{\rm keV}]$,[33] $m_{f_0}=0.969\,$GeV, $g^{(0)}_{\sigma\gamma\gamma}=-0.049\,{\rm GeV}^{-1}$ $[\Gamma^{(0)}_{\sigma\to\gamma\gamma}(m^2_\sigma)$ miserable], $g^{(0)}_{f_0\gamma\gamma}=0.718\,{\rm GeV}^{-1}$ $[\Gamma^{(0)}_{f_0\to\gamma\gamma}(m^2_{f_0})\approx0.01$ keV], $x_1=0.9$ GeV and $x_2=1.75$ GeV. It is clear from comparison of Figs. 17(b) and 18(b) that the form factor effect on the $\gamma\gamma\to\pi^0\pi^0$ cross section is weak in contrast to the $\gamma\gamma\to\pi^+\pi^-$ one [compare Figs. 17(a) and 18(a)] in which the σ_2 contribution is modified mainly. One emphasizes that all our conclusions about the mechanisms of the two-photon decays (productions) of the $\sigma(600)$ and $f_0(980)$ resonances are in force.[34]

---

[31] Such a natural modification of the point-like Born contribution was discussed in connection with the data on the $\gamma\gamma\to\pi^+\pi^-$ (Poppe, 1986; Morgan and Pennington, 1987, 1988; Boyer *et al.*, 1990; Behrend *et al.*, 1992) and $\gamma\gamma\to K^+K^-(K^0\bar K^0)$ reactions (Achasov and Shestakov, 1992, 1994b). But only the problem of the consistent description of the Belle data on $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ indicates the modification need of the Born sector of the model unambiguously (Achasov and Shestakov, 2008a, 2008b).

[32] At the same time we emphasize that the considerable systematic errors, the sources of which are described in detail in (Mori *et al.*, 2007a, 2007b; Uehara *et al.*, 2008), do not depreciate the role of the high statistics of the data, which allows to resolve the small local effects connected with the $f_0(980)$ resonance manifestation.

[33] Our $f_2(1270)$ two-photon width (3.83 keV) is more than the PDG fit (2.7 keV). A plausible reason of this difference is the fact that we calculate the background under the $f_2(1270)$ resonance theoretically. It is interesting that our similar treatment (Achasov and Shestakov, 1988) of the Crystal Ball data on the $\gamma\gamma\to\pi^0\eta$ reaction (Antreasyan *et al.*, 1986) gave for the two-photon width of the $a_2(1320)$ resonance the value 1.3 keV greater than the PDG average 1 keV keeping the gold $q\bar q$ model relation 25:9.

[34] Notice that the point-like ω and $a_2(1320)$ exchanges in the $\gamma\gamma\to\pi^0\pi^0$ and $\gamma\gamma\to\pi^+\pi^-$ amplitude, respectively, which give the contributions (mainly the S wave one) in the cross section, runaway with increasing the energy and comparable with the $f_2(1270)$ resonance contribution even in its energy region, are not observed experimentally. This was puzzled out in our paper (Achasov and Shestakov, 1992) with the $\gamma\gamma\to\pi^0\eta$ example. The proper reggezation of the point-like exchanges with the high spins reduces the dangerous contributions greatly. In addition, the cancelations between the ω and $h_1(1170)$ exchanges in $\gamma\gamma\to\pi^0\pi^0$ and the $a_2(1320)$ and $a_1(1260)$ exchanges in $\gamma\gamma\to\pi^+\pi^-$ take place. As to the ρ exchange in $\gamma\gamma\to\pi\pi$, its contribution is small for $g^2_{\rho\pi\gamma}\approx(1/9)g^2_{\omega\pi\gamma}$ and canceled ad-

E. Conclusion: Two-Photon Decays on the Nature of the $\sigma(600)$ and $f_0(980)$ mesons

Thus the physics of the two-photon decays of the light scalar mesons acquires the rather clear outline. The mechanism of their decays into $\gamma\gamma$ does not look like the mechanism of the classic tensor $q\bar{q}$ meson decays, which is the direct annihilation $q\bar{q}\to\gamma\gamma$. The light scalar meson decays into $\gamma\gamma$ are suppressed in comparison with the the tensor meson ones. They are caused by the rescattering mechanisms, i.e., by the four-quark transitions $\sigma(600)\to\pi^+\pi^-\to\gamma\gamma$, $f_0(980)\to K^+K^-\to\gamma\gamma$, and $a_0(980)\to K^+K^-\to\gamma\gamma$. Such a picture is suggested by experiment and supports the $q^2\bar{q}^2$ nature of the light scalars. It is significant that in the scalar meson case the longing for the exhaustive characteristic of their coupling with photons via the constant values $\Gamma_{0^{++}\to\gamma\gamma}(m^2_{0^{++}})$ by analogy with the tensor mesons cannot be realized for quite a number reasons.

First of all it is clear that when we deal with resonances accompanied by fundamental background, when two-photon decay widths change sharply in the resonance region for close inelastic thresholds, then there is no point in discussing the two-photon width in the resonance peak.

In this connection it is interesting to consider the cross section $\gamma\gamma\to\pi^+\pi^-$ caused only the resonance contributions, i.e.,

$$\sigma_{\rm res}(\gamma\gamma\to\pi^+\pi^-;s)$$
$$=[\rho_{\pi^+}(s)/(32\pi s)]\left|\widetilde{I}_{\pi^+\pi^-}(s;x_1)\,e^{2i\delta_B^{\pi\pi}(s)}T^{\pi\pi}_{\rm res}(s)\right.$$
$$\left.+\,\widetilde{I}_{K^+K^-}(s;x_2)\,T_{K^+K^-\to\pi^+\pi^-}(s)+M^{\rm direct}_{\rm res}(s)\right|^2 \quad (34)$$

[see Eqs. (10) and (25)–(27)], where the function $\widetilde{I}_{\pi^+\pi^-}(s;x_1)$ and $\widetilde{I}_{K^+K^-}(s;x_2)$ are the analogs of the $\widetilde{I}_{\pi^+\pi^-}(s)$ and $\widetilde{I}_{K^+K^-}(s)$ functions constructed with taking the form factors into account, see Appendixes A and C. Fig. 19 shows the dependence of the $\sigma_{\rm res}(\gamma\gamma\to\pi^+\pi^-;s)$ cross section, multiplied by the factor $3s/(8\pi^2)$, on the energy. In and around 1 GeV there is the impressive peak from the $f_0(980)$ resonance due to the inelastic $\gamma\gamma\to K^+K^-\to\pi^+\pi^-$ rescattering in the main. Following (Achasov and Shestakov, 1988, 2005, 2008b) one determines the $f_0(980)\to\gamma\gamma$ decay width averaged over the resonance mass distribution in the $\pi\pi$ channel

$$\langle\Gamma_{f_0\to\gamma\gamma}\rangle_{\pi\pi}=\int\limits_{0.8\ \rm GeV}^{1.1\ \rm GeV}\frac{3s}{8\pi^2}\sigma_{\rm res}(\gamma\gamma\to\pi^+\pi^-;s)d\sqrt{s}. \quad (35)$$

This value is the adequate functional characteristic of the coupling of the $f_0(980)$ with $\gamma\gamma$. For the presented combined fitting, $\langle\Gamma_{f_0\to\gamma\gamma}\rangle_{\pi\pi}\approx 0.19\,$keV (Achasov and Shestakov, 2008b). Taking into account that the wide $\sigma(600)$-resonance dominates in region $2m_\pi<\sqrt{s}<0.8\,$GeV one obtains by analogy with (35) $\langle\Gamma_{\sigma\to\gamma\gamma}\rangle_{\pi\pi}\approx 0.45\,$keV (Achasov and Shestakov, 2008b). Note that the cusp near the $\pi\pi$ threshold in the $[3s/(8\pi^2)]\sigma_{\rm res}(\gamma\gamma\to\pi^+\pi^-;s)$ expression, shown in Fig. 19, is the manifestation of the correction for the finite width in the propagator of the scalar resonance. In Appendix A there are adduced the transparent explanation of this phenomenon. In the total S wave amplitude $\gamma\gamma\to\pi\pi$ such a threshold enhancement is absent due to shielding the resonance contribution in the amplitude $T^0_0(s)$ by the chiral background one, see, for example, Fig. 18(b).



Figure 19 The integrand in Eq. (35) corresponding to the joint fit to the $\gamma\gamma\to\pi^+\pi^-$ and $\gamma\gamma\to\pi^0\pi^0$ data is shown by the solid curve. The dotted and dashed curves show the contributions from the resonant elastic $\gamma\gamma\to\pi^+\pi^-\to\pi^+\pi^-$ and inelastic $\gamma\gamma\to K^+K^-\to\pi^+\pi^-$ rescatterings, respectively.

The above examples, each in its manner give to feel clear the nontriviality of accessing information about the $\sigma(600)$ and $f_0(980)$ decays into $\gamma\gamma$. For instance, to determine $\Gamma_{\sigma\to\gamma\gamma}(m^2_\sigma)$ directly from data is impossible for the cross section in the σ region is formed by both resonance and compensating background. The specific dynamic model of the total amplitude needs to their separation. The simple Breit-Wigner is not enough here.

As for the $f_0(980)$ resonance, experimenters began to take into consideration two from three important circumstances (Mori *et al.*, 2007a, 2007b; Uehara *et al.*, 2008; see also Amsler *et al.*, 2008), for which we drew attention in (Achasov and Shestakov, 2005). Firstly, there was taken account the correction for the finite width due to the coupling of $f_0(980)$ with $K\bar{K}$ channel in the $f_0(980)$ resonance propagator, which effects essentially on the shape of the $f_0(980)$ peak in the $\pi\pi$ channel. Secondly, there was taken into account the interference of the $f_0(980)$ resonance with the background though in the simplest form. But no model was constructed for the $f_0(980)\to\gamma\gamma$ decay amplitude which was approximated simply by a constant (Mori *et al.*, 2007a, 2007b; Uehara *et al.*, 2008). Fitting data in this way the Belle collaboration extracted the values for $\Gamma_{f_0\to\gamma\gamma}(m^2_{f_0})$ presented in Table IV. But the discussion needs how to understand these values. First and foremost, one cannot use them for determining a

---

ditionally by the $b_1(1235)$ one.

coupling constant $g_{f_0\gamma\gamma}$ in a effective lagrangian, i.e., a constant of the direct transition $f_0(980)\to\gamma\gamma$, because such a constant is small and does not determine the $f_0(980)\to\gamma\gamma$ decay, as shown above. Until the model of the $f_0(980)\to\gamma\gamma$ decay amplitude is not specified, a meaning of the $\Gamma_{f_0\to\gamma\gamma}(m^2_{f_0})$ values, extracted with the help of the simplified parametrization, is rather vague.[35] In principle the $\Gamma_{f_0\to\gamma\gamma}(m^2_{f_0})$ values in Table IV can be taken as the preliminary estimations of $\langle\Gamma_{f_0\to\gamma\gamma}\rangle$, i.e., as the $f_0(980)\to\gamma\gamma$ decay width averaged over the hadron mass distribution (Achasov and Shestakov, 1988, 2005, 2008b).

In the dispersion approach there are introduced usually the pole two-photon widths $\Gamma_{R\to\gamma\gamma}(pole)$, $R=\sigma, f_0$ to characterize the coupling $\sigma(600)$ and $f_0(980)$ resonances with photons (see, for example, Morgan and Pennington, 1988, 1990; Boglione and Pennington, 1999, Pennington, 2006; Pennington, Mori, Uehara, and Watanabe, 2008). These widths are determined through the modulli of the complex pole residues of the $\gamma\gamma\to\pi\pi$ and $\pi\pi\to\pi\pi$ partial amplitudes constructed theoretically. Basing on our investigation (Achasov and Shestakov, 2007) we would like to note the following. The residues of the above amplitudes are essentially complex and cannot be used as any coupling constants in a hermitian effective lagrangian. These residues are "dressed" by the background for they relate to the total amplitudes. As our analysis in the $SU(2)_L\times SU(2)_R$ linear σ-model (Achasov and Shestakov, 2007) indicated, the background effects essentially on the values and phases of the residues. Thus the focus on the values of the $\Gamma_{R\to\gamma\gamma}(pole)$ type in dispersion approach does not help to reveal the mechanism of the two-photon decays of the scalar mesons and so cannot shed light on the nature of the light scalars.

IV. FUTURE TRENDS

A. The $a_0(980)$ Meson in the Reaction $\gamma\gamma\to\pi^0\eta$

Our conclusions about the important role of the K^+K^- loop mechanism in the two-photon production of the $a_0(980)$ resonance and its possible four-quark nature (Achasov and Shestakov, 1988, 1991, 1992) were based on the analysis of the results of the first experiments Crystal Ball (Antreasyan *et al.*, 1986; see Fig. 4) and JADE (Oest *et al.*, 1990) on the $\gamma\gamma\to\pi^0\eta$ reaction. Unfortunately, the large statistical errors in these data and the rather rough step of the $\pi^0\eta$ invariant mass distribution (equal 40 MeV in the Crystal Ball experiment and 60 MeV in the JADE one) left many uncertainties.



Figure 20 (a) Cross section for the reaction $\gamma\gamma\to\pi^0\eta$, without restriction on $|\cos\theta|$. The dashed, solid, and dotted curves correspond to the $a_2(1320)$ production, the $a_0(980)$ plus $a_2(1320)$ production, and the cross section folded (in the $a_0(980)$ region) with a Gaussian with $\sigma=10$ MeV mass resolution. The $a_0(980)$ is produced via the rescattering mechanism $\gamma\gamma\to K^+K^-\to a_0(980)\to\pi^0\eta$. The $a_2(1320)$ is produced via the direct coupling to photons; the parameterization of the $\gamma\gamma\to a_2(1320)\to\pi^0\eta$ cross section is the same as given by Antreasyan *et al.* (1986), but with the current values of the $a_2(1320)$ resonance parameters from Amsler *et al.* (2008). (b) The energy-dependent width of the $a_0(980)\to K^+K^-\to\gamma\gamma$ decay.

Both Crystal Ball and JADE registered events of production of $\pi^0\pi^0\to\gamma\gamma\gamma\gamma$ and $\pi^0\eta\to\gamma\gamma\gamma\gamma$ simultaneously. Because of this, we cherish the hope for appearing the Belle data on the $\gamma\gamma\to\pi^0\eta$ reaction in the $a_0(980)$ and $a_2(1320)$ resonances with the statistics exceeding the former data in a hundred times like the $\gamma\gamma\to\pi^0\pi^0$ Belle data. It is believed (see discussions in Achasov and Shestakov, 1988, 1992) that in the $\gamma\gamma\to\pi^0\eta$ reaction the background conditions in the $a_0(980)$ region are rather favorable. In particular, the non-coherent background in the $a_0(980)$ region from the $a_2(1320)$ resonance in $\gamma\gamma\to\pi^0\eta$ should be one order of magnitude smaller than the non-coherent one in the $f_0(980)$ region from the $f_2(1270)$ resonance in $\gamma\gamma\to\pi^0\pi^0$ [see, for example, Figs. 7(b) and 18(b)], because $[B(a_2\to\gamma\gamma)B(a_2\to\pi\eta)]/[B(f_2\to\gamma\gamma)B(f_2\to\pi^0\pi^0)]\times(\Gamma^2_{a_2}/\Gamma^2_{f_2})\approx 0.334\times 0.342=0.114$ (Amsler *at al.*, 2008). In Fig. 20(a) The reference theoretical curve for the $\gamma\gamma\to[a_0(980)+a_2(1320)]\to\pi^0\eta$ cross section is given, which corresponds to the $a_0(980)$ resonance production by the $\gamma\gamma\to K^+K^-\to a_0(980)\to\pi^0\eta$ rescattering and the $a_2(1320)$ resonance production by the direct coupling of $a_2(1320)$ with $\gamma\gamma$ (see Appendix B for details). In Fig. 20(b) the dependence of the $a_0(980)\to K^+K^-\to\gamma\gamma$ decay width on energy is shown. Averaging this over the resonance mass distribution in the $\pi^0\eta$ channel in the interval from 0.8 up 1.1 GeV , see Eq. (68) in Appendix B, one obtains $\langle\Gamma_{a_0\to K^+K^-\to\gamma\gamma}\rangle_{\pi^0\eta}\approx 0.125\,\text{keV}$.

B. The $f_0(980)$ and $a_0(980)$ Mesons near the $\gamma\gamma\to K\bar{K}$ Reaction Thresholds

The four-quark nature of the $a_0(980)$ and $f_0(980)$ resonances shows itself paradoxically in the $\gamma\gamma\to K\bar{K}$

---

[35] The above comments are true also in the $\sigma(600)$ resonance case.



Figure 21 The K^+K^- (a) and $K^0_SK^0_S$ (b) mass spectra measured by ARGUS (Albrecht *et al.*, 1990) and L3 (Acciarri *et al.*, 2001). (c) This plot illustrates the scale of the $K\bar{K}$ production cross section. The experimental points show the cross section for $\gamma\gamma\to K^+K^-$ with allowed contributions from $\lambda J=[22,\ 02,\ 00]$ (ARGUS). The upper dashed, dot-dashed, and solid curves correspond to the $\gamma\gamma\to K^+K^-$ Born cross section with $\lambda J=00$, $\lambda J=[00,\ 22]$, and the total one, respectively (the $\lambda J=02$ contribution is negligible). The lower dashed, dot-dashed, and solid curves correspond to the same cross sections but modified by a form factor (see Sec. III. D and Appendix A). The dotted curve shows the estimate of the the S wave $\gamma\gamma\to K^+K^-$ cross section in our model.

reactions (Achasov and Shestakov, 1992, 1994b). The $a_0(980)$ and $f_0(980)$ production is suppressed for the strong final state interaction in the $\gamma\gamma\to K^+K^-$ reaction and in the $\gamma\gamma\to K^0K^0$ one for the destructive the $a_0(980)$ and $f_0(980)$ interference.

A few words about the experimental situation. The $\gamma\gamma\to K^+K^-$ and $\gamma\gamma\to K^0_SK^0_S$ cross sections become saturated in fact with the contributions of the $f_2(1270)$, $a_2(1320)$, and $f_2'(1525)$ resonances being produced in the $\lambda=2$ helicity states in the energy region $1<\sqrt{s}<1.7$ GeV (Althoff *et al.*, 1983, 1985; Johnson, 1986; Berger *et al.*, 1988; Behrend *et al.*, 1989; Albrecht *et al.*, 1990; Feindt and Harjes, 1990; Braccini, 2000; Acciarri *et al.*, 2001; Abe *et al.*, 2004), see Figs. 21(a) and 21(b). The constructive and destructive interference between the $f_2(1270)$ and $a_2(1320)$ resonance contributions is observed in $\gamma\gamma\to K^+K^-$ and $\gamma\gamma\to K^0\bar{K}^0$, respectively, in agreement with the $q\bar{q}$ model (Faiman, Lipkin, and Rubinstein, 1975). Notice that the region of the $K\bar{K}$ thresholds, $2m_K<\sqrt{s}<1.1$ GeV, sensitive to the S wave contributions is not investigated in fact. The sensitivity of the ARGUS experiment to the K^+K^- events at $2m_{K^+}<\sqrt{s}<1.1$ GeV was negligible (Albrecht *et al.*, 1990), see Fig. 21(a), and the total statistics of the L3 (Acciarri *et al.*, 2001), see Fig. 21(b), and CLEO (Braccini, 2000) experiments on $\gamma\gamma\to K^0_SK^0_S$ at $2m_{K^0}<\sqrt{s}<1.1$ GeV are within 60 events.

The absence of the considerable non-resonance background in the $\gamma\gamma\to K^+K^-$ cross section seems at first sight rather surprising for the Born contribution comparable with the tensor resonance contributions, see Fig. 21(c). As seen from this figure, the S wave contribution dominates in the Born cross section at $\sqrt{s}<1.5$ GeV. One would think that the large non-coherent background should be under the tensor meson peaks in the K^+K^- channel. But taking account of the resonance interaction between the K^+ and K^- mesons in the final state results in the cancelation of the considerable part of this background (Achasov and Shestakov, 1992, 1994b). The principal point is that the Born S wave $\gamma\gamma\to K^+K^-$ amplitude acquires the $\xi(s)=[1+i\rho_{K^+}(s)T_{K^+K^-\to K^+K^-}(s)]$ factor due to the $\gamma\gamma\to K^+K^-\to K^+K^-$ rescattering amplitude with the real kaons in the intermediate state. The $a_0(980)$ and $f_0(980)$ resonance contributions dominate in the $T_{K^+K^-\to K^+K^-}(s)$ amplitude near the K^+K^- threshold and provide it with the considerable imaginary part for the strong coupling with the $K\bar{K}$ channels in the four-quark scheme. As a result the $|\xi(s)|^2$ factor is considerably less than 1 just above the K^+K^- threshold and the seed S wave Born contribution is considerably reduced in the wide region of $\sqrt{s}$. The dotted curve in Fig. 21(c) represents the estimation of the S wave $\gamma\gamma\to K^+K^-$ cross section obtained in the model under consideration with taking the form factor into account (see details in Appendix C). This estimation agrees with obtained earlier (Achasov and Shestakov, 1992, 1994b).

So one can hope to detect in the partial wave analysis of the $\gamma\gamma\to K^+K^-$ reaction at $2m_{K^+}<\sqrt{s}<1.1$ GeV the scalar contributions at the rate of about 5–10 nb. As for the $\gamma\gamma\to K^0\bar{K}^0$ reaction, its amplitude has not the Born contribution and the $a_0(980)$ resonance contribution has the opposite sign in comparison with the $\gamma\gamma\to K^+K^-$ channel. As a result the contributions of the S wave $\gamma\gamma\to K^+K^-\to K^0\bar{K}^0$ rescattering amplitudes with the isotopic spin $I=0$ and 1 practically cancel each other and the the corresponding cross section should be at the rate of about $\lesssim 1$ nb.

Acknowledgments

This work was supported in part by the RFFI Grant No. 07-02-00093 from the Russian Foundation for Basic Research and by the Presidential Grant No. NSh-1027.2008.2 for Leading Scientific Schools.

V. APPENDIX

A. $\gamma\gamma\to\pi\pi$

Below there is the list of the expressions for the Born helicity amplitudes $M^{\rm Born}_\lambda(s,\theta)$, the loop integrals $\widetilde{I}_{\pi^+\pi^-}(s)$ and $\widetilde{I}_{K^+K^-}(s)$ from Eqs. (10) – (13), some low partial Born amplitudes, $M^{\rm Born}_{\lambda J}(s)$, and the Born partial amplitudes with taking into account the form factor. In addition, a few useful auxiliary formulae for the solitary scalar resonance are adduced.

The Born helicity $\gamma\gamma\to\pi^+\pi^-$ amplitudes are

$$M^{\rm Born}_0(s,\theta)=\frac{4m^2_{\pi^+}}{s}\,\frac{8\pi\alpha}{1-\rho^2_{\pi^+}(s)\cos^2\theta}\,,$$

$$\rho_{\pi^+}(s)=(1-4m^2_{\pi^+}/s)^{1/2}\,, \qquad (36)$$

$$M_2^{\rm Born}(s,\theta)=\frac{8\pi\alpha\rho^2_{\pi^+}(s)\sin^2\theta}{1-\rho^2_{\pi^+}(s)\cos^2\theta} \qquad (37)$$

($\alpha=e^2/(4\pi)=1/137$). Their partial wave series expansion is

$$M_\lambda^{\rm Born}(s,\theta)=\sum_{J\geq\lambda}(2J+1)M_{\lambda J}^{\rm Born}(s)d^J_{\lambda 0}(\theta)\,, \qquad (38)$$

where $d^J_{\lambda 0}(\theta)$ are the usual d functions (see, for example, Amsler *et al.*, 2008). Three low partial waves are

$$M_{00}^{\rm Born}(s)=4\pi\alpha\frac{1-\rho^2_{\pi^+}(s)}{\rho_{\pi^+}(s)}\ln\frac{1+\rho_{\pi^+}(s)}{1-\rho_{\pi^+}(s)}\,, \qquad (39)$$

$$M_{02}^{\rm Born}(s)=4\pi\alpha\frac{1-\rho^2_{\pi^+}(s)}{\rho^2_{\pi^+}(s)}\left[\frac{3-\rho^2_{\pi^+}(s)}{2\rho_{\pi^+}(s)}\right.$$

$$\left.\times\ln\frac{1+\rho_{\pi^+}(s)}{1-\rho_{\pi^+}(s)}-3\right]\,, \qquad (40)$$

$$M_{22}^{\rm Born}(s)=4\pi\alpha\sqrt{\frac{3}{2}}\left[\frac{(1-\rho^2_{\pi^+}(s))^2}{2\rho^3_{\pi^+}(s)}\ln\frac{1+\rho_{\pi^+}(s)}{1-\rho_{\pi^+}(s)}\right.$$

$$\left.-\frac{1}{\rho^2_{\pi^+}(s)}+\frac{5}{3}\right]\,. \qquad (41)$$

The function $\widetilde{I}_{\pi^+\pi^-}(s)$ is the amplitude of the triangle loop diagram $\gamma\gamma\to\pi^+\pi^-\to$ *a scalar with the mass* $=\sqrt{s}$,

$$\widetilde{I}_{\pi^+\pi^-}(s)=\frac{s}{\pi}\int_{4m^2_{\pi^+}}^\infty\frac{\rho_{\pi^+}(s')M_{00}^{\rm Born}(s')}{s'(s'-s-i\varepsilon)}ds'\,. \qquad (42)$$

The behavior $\widetilde{I}_{\pi^+\pi^-}(s)\propto s$ when $s\to 0$, is the gauge invariance consequence. At $s\geq 4m^2_{\pi^+}$

$$\widetilde{I}_{\pi^+\pi^-}(s)=8\alpha\left\{\frac{m^2_{\pi^+}}{s}\left[\pi+i\ln\frac{1+\rho_{\pi^+}(s)}{1-\rho_{\pi^+}(s)}\right]^2-1\right\}\,. \qquad (43)$$

The form factor, see Eqs. (33),

$$G(t,u)=\frac{1}{s}\left[\frac{m^2_{\pi^+}-t}{1-(u-m^2_{\pi^+})/x_1^2}+\frac{m^2_{\pi^+}-u}{1-(t-m^2_{\pi^+})/x_1^2}\right]\}$$

(where $t=m^2_{\pi^+}-s[1-\rho_{\pi^+}(s)\cos\theta]/2$ and $u=m^2_{\pi^+}-s[1+\rho_{\pi^+}(s)\cos\theta]/2$) modifies the Born partial wave amplitudes. Let us introduce the notations

$$M_{0J}^{\rm Born}(s)=[(1-\rho^2_{\pi^+}(s))/\rho_{\pi^+}(s)]F_{0J}^{\rm Born}(\rho_{\pi^+}(s))\,, \qquad (44)$$

$$M_{2J}^{\rm Born}(s)=\rho_{\pi^+}(s)F_{2J}^{\rm Born}(\rho_{\pi^+}(s))\,. \qquad (45)$$

Then the modified amplitudes can be represented in the form

$$M_{0J}^{\rm Born}(s;x_1)$$

$$=\frac{1-\rho^2_{\pi^+}(s)}{\rho_{\pi^+}(s)}\left[F_{0J}^{\rm Born}(\rho_{\pi^+}(s))-F_{0J}^{\rm Born}(\rho_{\pi^+}(s;x_1))\right]\,, \qquad (46)$$

$$M_{2J}^{\rm Born}(s;x_1)$$

$$=\rho_{\pi^+}(s)\left[F_{2J}^{\rm Born}(\rho_{\pi^+}(s))-F_{2J}^{\rm Born}(\rho_{\pi^+}(s;x_1))\right]\,, \qquad (47)$$

where

$$\rho_{\pi^+}(s;x_1)=\rho_{\pi^+}(s)/(1+2x_1^2/s)\,. \qquad (48)$$

As to the function $\widetilde{I}_{\pi^+\pi^-}(s)$, see (42) and (43), this is replaced by

$$\widetilde{I}_{\pi^+\pi^-}(s;x_1)=\frac{s}{\pi}\int_{4m^2_{\pi^+}}^\infty\frac{\rho_{\pi^+}(s')M_{00}^{\rm Born}(s';x_1)}{s'(s'-s-i\varepsilon)}ds'\,. \qquad (49)$$

In this case the numerical integration needs certainly.

To make easy understanding the structure and normalization of the sufficiently complicated expressions used in fitting data, one adduces the formulae of the σ production due to the $\gamma\gamma\to\pi^+\pi^-\to\sigma\to\pi^+\pi^-$ rescattering and its two-photon decay in the imaginary case of the solitary scalar σ resonance coupled only to the $\pi\pi$ channel.

The corresponding resonance cross section has the familiar form

$$\sigma_{\rm res}(\gamma\gamma\to\pi^+\pi^-)$$

$$=\frac{8\pi}{s}\,\frac{\sqrt{s}\Gamma_{\sigma\to\pi^+\pi^-\to\gamma\gamma}(s)\,\sqrt{s}\Gamma_{\sigma\to\pi^+\pi^-}(s)}{|D_\sigma(s)|^2}\,, \qquad (50)$$

where

$$\Gamma_{\sigma\to\pi^+\pi^-\to\gamma\gamma}(s)=\frac{1}{16\pi\sqrt{s}}|M_{\sigma\to\pi^+\pi^-\to\gamma\gamma}(s)|^2$$

$$=\left|\frac{1}{16\pi}\,\widetilde{I}_{\pi^+\pi^-}(s)\right|^2\frac{g^2_{\sigma\pi^+\pi^-}}{16\pi\sqrt{s}}\,. \qquad (51)$$

If σ can else directly transit into $\gamma\gamma$ with the amplitude $sg^{(0)}_{\sigma\gamma\gamma}$ then the width $\Gamma_{\sigma\to\pi^+\pi^-\to\gamma\gamma}(s)$ in (50) should be replaced by

$$\Gamma_{\sigma\to\gamma\gamma}(s)=\frac{1}{16\pi\sqrt{s}}|M_{\sigma\to\gamma\gamma}(s)|^2\,, \qquad (52)$$

where

$$M_{\sigma\to\gamma\gamma}(s)=M_{\sigma\to\pi^+\pi^-\to\gamma\gamma}(s)+sg^{(0)}_{\sigma\gamma\gamma}\,. \qquad (53)$$

The propagator of the σ resonance with the m_σ Breit-Wigner mass in (50) has the form

$$\frac{1}{D_\sigma(s)}=\frac{1}{m^2_\sigma-s+{\rm Re}\Pi^{\pi\pi}_\sigma(m^2_\sigma)-\Pi^{\pi\pi}_\sigma(s)}\,, \qquad (54)$$



Figure 22 Demonstration of the finite width correction with an example of the single σ resonance. The curves are described in the text.

where $\Pi^{\pi\pi}_\sigma(s)$ is the polarization operator of the σ resonance for the contribution of the $\pi^+\pi^-$ and $\pi^0\pi^0$ intermediate states. At $s\geq 4m^2_{\pi^+}\,(=4m^2_{\pi^0})$

$$\Pi^{\pi\pi}_\sigma(s)=\frac{3}{2}\frac{g^2_{\sigma\pi^+\pi^-}}{16\pi}\rho_{\pi^+}(s)\left[i-\frac{1}{\pi}\ln\frac{1+\rho_{\pi^+}(s)}{1-\rho_{\pi^+}(s)}\right]\,,\quad(55)$$

when $0<s<4m^2_{\pi^+}$ then $\rho_{\pi^+}(s)\to i|\rho_{\pi^+}(s)|$. The $\sigma\to\pi\pi$ decay width

$$\Gamma_{\sigma\to\pi\pi}(s)=\frac{1}{\sqrt{s}}\mathrm{Im}\Pi^{\pi\pi}_\sigma(s)=\frac{3}{2}\frac{g^2_{\sigma\pi^+\pi^-}}{16\pi}\frac{\rho_{\pi^+}(s)}{\sqrt{s}}\,.\quad(56)$$

The function $\mathrm{Re}[\Pi^{\pi\pi}_\sigma(m^2_\sigma)-\Pi^{\pi\pi}_\sigma(s)]$ in the denominator of Eq. (54) is the correction for the finite width of resonance. In Fig. 20 the real and imaginary parts[36] of the inverse propagator $D_\sigma(s)$ are shown by the solid and dashed curves in the case of the resonance with the mass $m_\sigma=0.6$ GeV and the width $\Gamma_\sigma=\Gamma_{\sigma\to\pi\pi}(m^2_\sigma)=0.45$ GeV. As may be inferred from this figure, $\mathrm{Re}[D_\sigma(s)]$ can be close to 0 at $s=4m^2_{\pi^+}$ due to the correction for the finite width in the case of the large one. Then this results in the threshold cusp in amplitudes $\propto|1/D_\sigma(s)|$.[37] For reference, in Fig. 20 the real and imaginary parts[38] of the inverse propagator $D_\sigma(s)=m^2_\sigma-s-i\Gamma_\sigma\sqrt{(s-4m^2_{\pi^+})/(m^2_\sigma-4m^2_{\pi^+})}$, in which the correction for the finite width is absent (Flatte, 1976), are shown by the dotted and dashed-dotted curves, respectively, at the same values of m_σ and Γ_σ.

B. $\gamma\gamma\to\pi^0\eta$

Below are the formulae which were used in graphing the curves in Fig. 20.

The helicity amplitudes $M_0(\gamma\gamma\to\pi^0\eta;s,\theta)$ and $M_2(\gamma\gamma\to\pi^0\eta;s,\theta)$, caused by the $a_0(980)$ and $a_2(1320)$ resonance production respectively, are

$$\begin{aligned}M_0(\gamma\gamma\to\pi^0\eta;s,\theta)&=M_{00}(\gamma\gamma\to\pi^0\pi^0;s)\\&=\widetilde{I}_{K^+K^-}(s;x_2)\,T_{K^+K^-\to a_0\to\pi^0\eta}(s),\end{aligned}\quad(57)$$

$$T_{K^+K^-\to a_0\to\pi^0\eta}(s)=\frac{g_{a_0K^+K^-}g_{a_0\pi\eta}}{16\pi}\frac{1}{D_{a_0}(s)},\quad(58)$$

$$\begin{aligned}M_2(\gamma\gamma\to\pi^0\eta;s,\theta)&=5d^2_{20}(\theta)M_{22}(\gamma\gamma\to\pi^0\eta;s)\\&=80\pi d^2_{20}(\theta)M_{\gamma\gamma\to a_2(1270)\to\pi^0\eta}(s).\end{aligned}\quad(59)$$

Their normalizations, the connections with the cross sections, are similar to the ones in Eqs. (16), (18), (28) and (31). The triangle loop integral $\widetilde{I}_{K^+K^-}(s;x_2)$ is defined in Appendix C, Eq. (83); $1/D_{a_0}(s)$ is the $a_0(980)$ resonance propagator.

The integral cross section of the $\gamma\gamma\to\pi^0\eta$ reaction is

$$\begin{aligned}\sigma(\gamma\gamma\to\pi^0\eta)&=\sigma(\gamma\gamma\to a_0(980)\to\pi^0\eta;s)\\&+\sigma(\gamma\gamma\to a_2(1320)\to\pi^0\eta;s).\end{aligned}\quad(60)$$

Here

$$\begin{aligned}&\sigma(\gamma\gamma\to a_0(980)\to\pi^0\eta;s)\\&=8\pi\,\Gamma_{a_0\to K^+K^-\to\gamma\gamma}(s)\Gamma_{a_0\to\pi\eta}(s)/|D_{a_0}(s)|^2,\end{aligned}\quad(61)$$

$$\Gamma_{a_0\to\pi\eta}(s)=\frac{g^2_{a_0\to\pi\eta}}{16\pi}\frac{\rho_{\pi\eta}(s)}{\sqrt{s}}\,,\quad(62)$$

where $\rho_{\pi\eta}(s)=\sqrt{(1-m^2_+/s)(1-m^2_-/s)}$ and $m_\pm=m_\eta\pm m_\pi$. The $a_0(980)$ resonance propagator is

$$\frac{1}{D_{a_0}(s)}=\frac{1}{m^2_{a_0}-s+\sum_{ab}\mathrm{Re}\Pi^{ab}_{a_0}(m^2_{a_0})-\Pi^{ab}_{a_0}(s)}\,,\quad(63)$$

where $ab=\pi\eta$, K^+K^-, $K^0\bar{K}^0$. At $s\geq m^2_+$

$$\begin{aligned}\Pi^{\pi\eta}_{a_0}(s)=\frac{g^2_{a_0\to\pi\eta}}{16\pi}\Bigg[&\frac{m_+m_-}{\pi s}\ln\frac{m_\pi}{m_\eta}\\&+\rho_{\pi\eta}(s)\left(i-\frac{1}{\pi}\ln\frac{\sqrt{s-m^2_-}+\sqrt{s-m^2_+}}{\sqrt{s-m^2_-}-\sqrt{s-m^2_+}}\right)\Bigg]\,.\end{aligned}\quad(64)$$

At $s\geq 4m^2_{K^+}$

$$\begin{aligned}&\Pi^{K^+K^-}_{a_0}(s)=\frac{g^2_{a_0K^+K^-}}{16\pi}\rho_{K^+}(s)\left[i-\frac{1}{\pi}\ln\frac{1+\rho_{K^+}(s)}{1-\rho_{K^+}(s)}\right],\\&\rho_{K^+}(s)=(1-4m^2_{K^+}/s)^{1/2}\,.\end{aligned}\quad(65)$$

---

[36] (taken with the sign minus)

[37] The references to papers, in which the corrections for the finite widths and the analytic properties of the propagators of the realistic $f_0(980)$, $a_0(980)$, and $\sigma(600)$ resonances, are pointed out in Section II. In connection with the $\gamma\gamma\to\pi^0\eta$ and $\gamma\gamma\to\pi\pi$ reaction these corrections are discussed also in the papers (Achasov and Shestakov, 1988, 2005).

[38] (also taken with the sign minus)

When $0<s<4m^2_{K^+}$, then $\rho_{K^+}(s)\to i|\rho_{K^+}(s)|$ and

$$\Pi_{a_0}^{K^+K^-}(s)=-\frac{g^2_{a_0K^+K^-}}{16\pi}|\rho_{K^+}(s)|\left[1-\frac{1}{\pi}\arctan|\rho_{K^+}(s)|\right].\tag{66}$$

$\Pi_{a_0}^{K^0\bar K^0}(s)$ results from $\Pi_{a_0}^{K^+K^-}(s)$ by the substitution of $K^0(\bar K^0)$ for $K^+(K^-)$.

The $a_0\to K^+K^-\to\gamma\gamma$ decay width in Eq. (61)

$$\Gamma_{a_0\to K^+K^-\to\gamma\gamma}(s)=\left|\frac{1}{16\pi}\,\widetilde{I}_{K^+K^-}(s;x_2)\right|^2\frac{g^2_{a_0\to K^+K^-}}{16\pi\sqrt{s}}\,.\tag{67}$$

The dependence of the $\Gamma_{a_0\to K^+K^-\to\gamma\gamma}(s)$ width on $\sqrt{s}$ is shown in Fig. 20(b) and the its average was calculated in the following way

$$\langle\Gamma_{a_0\to K^+K^-\to\gamma\gamma}\rangle_{\pi\eta}$$

$$=\int\limits_{0.8\ \text{GeV}}^{1.1\ \text{GeV}}\Gamma_{a_0\to K^+K^-\to\gamma\gamma}(s)\frac{\sqrt{s}\Gamma_{a_0\to\pi\eta}(s)}{\pi|D_{a_0}(s)|^2}\,2\sqrt{s}\,d\sqrt{s}$$

$$=\int\limits_{0.8\ \text{GeV}}^{1.1\ \text{GeV}}\frac{s}{4\pi^2}\sigma(\gamma\gamma\to a_0(980)\to\pi^0\eta;s)\,d\sqrt{s}.\tag{68}$$

We put $m_{a_0}=0.98\,\text{GeV}$, $g^2_{a_0^0K^+K^-}/(4\pi)=1.5\,\text{GeV}^2$ and used the naive $q^2\bar q^2$ model relation $g_{a_0\pi\eta}=\sqrt{2/3}\,g_{a_0^0K^+K^-}$ (Achasov and Shestakov, 1988, 1992; Achasov and Kiselev, 2006, 2007a).

For the $\gamma\gamma\to a_2(1320)\to\pi^0\eta$ cross section in Eq. (60) we used the parametrization from (Antreasyan *et al.*, 1986)

$$\sigma(\gamma\gamma\to a_2(1320)\to\pi^0\eta;s)$$

$$=\frac{40\pi\,\Gamma_{a_2\to\gamma\gamma}B(a_2\to\pi\eta)\Gamma^{\text{tot}}_{a_2}(s)}{(m^2_{a_2}-s)^2+m^2_{a_2}\Gamma^{\text{tot}}_{a_2}(s)^2}\tag{69}$$

and the current values of the $a_2(1320)$ resonance parameters, which are adduced by PDG (Amsler *et al.*, 2008).

The energy dependence of the total width in Eq. (69) is approximated in the following way

$$\Gamma^{\text{tot}}_{a_2}(s)=\Gamma^{\text{tot}}_{a_2}(m^2_{a_2})\frac{q^5_{\pi\eta}(s)}{q^5_{\pi\eta}(m^2_{a_2})}\frac{D_2(q_{\pi\eta}(m^2_{a_2})R_{a_2})}{D_2(q_{\pi\eta}(s)R_{a_2})}\,,\tag{70}$$

where $q_{\pi\eta}(s)=\sqrt{s}\rho_{\pi\eta}(s)/2$, $D_2(x)=9+3x^2+x^4$ and $R_{a_2}=5.07\ \text{GeV}^{-1}$ (or 1 fm).

C. $\gamma\gamma\to K\bar K$

The $\gamma\gamma\to K^+K^-$ Born amplitudes $\widehat{M}^{\text{Born}}_{\lambda}(s,\theta)$ and $\widehat{M}^{\text{Born}}_{\lambda J}(s)$ result from the $\gamma\gamma\to\pi^+\pi^-$ Born amplitudes $M^{\text{Born}}_{\lambda}(s,\theta)$ and $M^{\text{Born}}_{\lambda J}(s)$ by the substitution of K^+ for π^+ in Eqs. (36), (37) and (39) – (41)

$$\widehat{M}^{\text{Born}}_{0}(s,\theta)=\frac{4m^2_{K^+}}{s}\frac{8\pi\alpha}{1-\rho^2_{K^+}(s)\cos^2\theta}\,,$$

$$\rho_{K^+}(s)=(1-4m^2_{K^+}/s)^{1/2}\,,\tag{71}$$

$$\widehat{M}^{\text{Born}}_{2}(s,\theta)=\frac{8\pi\alpha\rho^2_{K^+}(s)\sin^2\theta}{1-\rho^2_{K^+}(s)\cos^2\theta}\,,\tag{72}$$

$$\widehat{M}^{\text{Born}}_{00}(s)=4\pi\alpha\frac{1-\rho^2_{K^+}(s)}{\rho_{K^+}(s)}\ln\frac{1+\rho_{K^+}(s)}{1-\rho_{K^+}(s)}\,,\tag{73}$$

$$\widehat{M}^{\text{Born}}_{02}(s)=4\pi\alpha\frac{1-\rho^2_{K^+}(s)}{\rho^2_{K^+}(s)}\left[\frac{3-\rho^2_{K^+}(s)}{2\rho_{K^+}(s)}\right.$$

$$\left.\times\ln\frac{1+\rho_{K^+}(s)}{1-\rho_{K^+}(s)}-3\right]\,,\tag{74}$$

$$\widehat{M}^{\text{Born}}_{22}(s)=4\pi\alpha\sqrt{\frac{3}{2}}\left[\frac{(1-\rho^2_{K^+}(s))^2}{2\rho^3_{K^+}(s)}\ln\frac{1+\rho_{K^+}(s)}{1-\rho_{K^+}(s)}\right.$$

$$\left.-\frac{1}{\rho^2_{K^+}(s)}+\frac{5}{3}\right]\,.\tag{75}$$

At $s\geq 4m^2_{K^+}$ the triangle loop function $\widetilde{I}_{K^+K^-}(s)$ results from $\widetilde{I}_{\pi^+\pi^-}(s)$ by the substitution of K^+ for π^+ in Eq. (43)

$$\widetilde{I}_{K^+K^-}(s)=8\alpha\left\{\frac{m^2_{K^+}}{s}\left[\pi+i\ln\frac{1+\rho_{K^+}(s)}{1-\rho_{K^+}(s)}\right]^2-1\right\}.\tag{76}$$

At $0<s<4m^2_{K^+}$ the analytic continuation results in

$$\widetilde{I}_{K^+K^-}(s)=8\alpha\left\{\frac{m^2_{K^+}}{s}\left[\pi-2\arctan|\rho_{K^+}(s)|\right]^2-1\right\}.\tag{77}$$

The

$$\widehat{G}(t,u)=\frac{1}{s}\left[\frac{m^2_{K^+}-t}{1-(u-m^2_{K^+})/x_2^2}+\frac{m^2_{K^+}-u}{1-(t-m^2_{K^+})/x_2^2}\right]\tag{78}$$

(here $t=m^2_{K^+}-s[1-\rho_{K^+}(s)\cos\theta]/2$ and $u=m^2_{K^+}-s[1+\rho_{K^+}(s)\cos\theta]/2$) modifies the partial waves amplitudes. The substitution of K^+ for π^+ in Eqs. (44) and (45) gives the auxiliary representation of $\widehat{M}^{\text{Born}}_{\lambda J}(s)$

$$\widehat{M}^{\text{Born}}_{0J}(s)=[(1-\rho^2_{K^+}(s))/\rho_{K^+}(s)]\widehat{F}^{\text{Born}}_{0J}(\rho_{K^+}(s)),\tag{79}$$

$$\widehat{M}^{\text{Born}}_{2J}(s)=\rho_{K^+}(s)\widehat{F}^{\text{Born}}_{2J}(\rho_{K^+}(s)),\tag{80}$$

The modified partial wave amplitudes result from Eqs. (46) and (47) by the substitution of K^+ and x_2 for π^+ and x_1.

$$\widehat{M}^{\rm Born}_{0J}(s;x_2) = \frac{1-\rho^2_{K^+}(s)}{\rho_{K^+}(s)}\Big[\widehat{F}^{\rm Born}_{0J}(\rho_{K^+}(s)) - \widehat{F}^{\rm Born}_{0J}(\rho_{K^+}(s;x_2))\Big], \quad (81)$$

$$\widehat{M}^{\rm Born}_{2J}(s;x_2) = \rho_{K^+}(s)\Big[\widehat{F}^{\rm Born}_{2J}(\rho_{K^+}(s)) - \widehat{F}^{\rm Born}_{2J}(\rho_{K^+}(s;x_2))\Big]. \quad (82)$$

The triangle loop function $\widetilde{I}_{K^+K^-}(s)$, modified by the form factor, is replaced by

$$\widetilde{I}_{K^+K^-}(s;x_2) = \frac{s}{\pi}\int\limits_{4m^2_{K^+}}^{\infty}\frac{\rho_{K^+}(s')\widehat{M}^{\rm Born}_{00}(s';x_2)}{s'(s'-s-i\varepsilon)}ds'. \quad (83)$$

The S wave amplitudes of the $\gamma\gamma\to K^+K^-$ and $\gamma\gamma\to K^0\bar{K}^0$ reactions in our model are

$$M_{00}(\gamma\gamma\to K^+K^-;s) = \widehat{M}^{\rm Born}_{00}(s;x_2) + \widetilde{I}_{\pi^+\pi^-}(s;x_1)T_{\pi^+\pi^-\to K^+K^-}(s) + \widetilde{I}_{K^+K^-}(s;x_2) \times T_{K^+K^-\to K^+K^-}(s) + M^{\rm direct}_{{\rm res};+}(s), \quad (84)$$

$$M_{00}(\gamma\gamma\to K^0\bar{K}^0;s) = \widetilde{I}_{\pi^+\pi^-}(s;x_1)\,T_{\pi^+\pi^-\to K^0\bar{K}^0}(s) + \widetilde{I}_{K^+K^-}(s;x_2) \times T_{K^+K^-\to K^0\bar{K}^0}(s) + M^{\rm direct}_{{\rm res};-}(s). \quad (85)$$

The corresponding cross sections are given by

$$\sigma_{00}(\gamma\gamma\to K^+K^-) = \frac{\rho_{K^+}(s)}{32\pi s}|M_{00}(\gamma\gamma\to K^+K^-;s)|^2 \quad (86)$$

and

$$\sigma_{00}(\gamma\gamma\to K^0_SK^0_S) = \frac{\rho_{K^0}(s)}{64\pi s}|M_{00}(\gamma\gamma\to K^0\bar{K}^0;s)|^2, \quad (87)$$

$\rho_{K^0}(s) = (1-4m^2_{K^0}/s)^{1/2}$. The amplitudes $\pi\pi\to K\bar{K}$, $T_{\pi^+\pi^-\to K^+K^-}(s) = T_{\pi^+\pi^-\to K^0\bar{K}^0}(s) = T_{K^+K^-\to\pi^+\pi^-}(s)$, are defined by Eqs. (23) and (26). The $K^+K^-\to K^+K^-$ and $K^+K^-\to K^0\bar{K}^0$ amplitudes are

$$T_{K^+K^-\to K^+K^-}(s) = [t^0_0(s)+t^1_0(s)]/2, \quad (88)$$

$$T_{K^+K^-\to K^0\bar{K}^0}(s) = [t^0_0(s)-t^1_0(s)]/2, \quad (89)$$

where $t^I_0(s)$ is the S wave amplitude of the $K\bar{K}\to K\bar{K}$ reaction with the isospin $I=0,1$;

$$t^0_0(s) = \frac{\exp[2i\delta^{K\bar{K}}_B(s)]-1}{2i\rho_{K^+}(s)} + e^{2i\delta^{K\bar{K}}_B(s)}T^{K\bar{K}}_{\rm res}(s)\,, \quad (90)$$

$$t^1_0(s) = g^2_{a_0K^+K^-}/[8\pi D_{a_0}(s)]\,, \quad (91)$$

$$T^{K\bar{K}}_{\rm res}(s) = \frac{g_{\sigma K^+K^-}\overline{\Delta}_{f_0}(s) + g_{f_0K^+K^-}\overline{\Delta}_{\sigma}(s)}{8\pi[D_\sigma(s)D_{f_0}(s)-\Pi^2_{f_0\sigma}(s)]}\,, \quad (92)$$

where $\overline{\Delta}_{f_0}(s) = D_{f_0}(s)g_{\sigma K^+K^-} + \Pi_{f_0\sigma}(s)g_{f_0K^+K^-}$ and $\overline{\Delta}_{\sigma}(s) = D_{\sigma}(s)g_{f_0K^+K^-} + \Pi_{f_0\sigma}(s)g_{\sigma K^+K^-}$. The amplitudes of the direct resonance transitions into photons

$$M^{\rm direct}_{{\rm res};\pm}(s) = s\,e^{i\delta^{K\bar{K}}_B(s)} \times \frac{g^{(0)}_{\sigma\gamma\gamma}\overline{\Delta}_{f_0}(s) + g^{(0)}_{f_0\gamma\gamma}\overline{\Delta}_{\sigma}(s)}{D_\sigma(s)D_{f_0}(s)-\Pi^2_{f_0\sigma}(s)} \pm s\frac{g^{(0)}_{a_0\gamma\gamma}g_{a_0K^+K^-}}{D_{a_0}(s)}\,, \quad (93)$$

where $g^{(0)}_{a_0\gamma\gamma}$ is the direct coupling constant of the $a_0(980)$ with $\gamma\gamma$. When estimating the S wave $\gamma\gamma\to K^+K^-$ cross section, see the dotted curve in Fig. 21(c), the amplitudes of the direct resonance transitions into photons were disregarded.

References

[1] Abarbanel, H.D.I., and M.L. Goldberger, 1968, Phys. Rev. **165**, 1594.
[2] Abe, K., *et al.* (Belle), 2004, Eur. Phys. J. C **32**, 323.
[3] Abe, K., *et al.* (Belle), 2007, BELLE-CONF-0768, eprint arXiv:0711.1926 [hep-ex].
[4] Acciarri,M., *et al.* (L3), 2001. Phys. Lett. B **501**, 173.
[5] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1979, Phys. Lett. B **88**, 367.
[6] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1980a, Sov. J. Nucl. Phys. **32**, 566 [Yad. Fiz. **32**, 1098 (1980)].
[7] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1980b, Phys. Lett. B **96**, 168.
[8] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1981a, Phys. Lett. B **102**, 196.
[9] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1981b, Sov. J. Nucl. Phys. **33**, 715 [Yad. Fiz. **33**, 1337 (1981)].
[10] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1981c, Preprint TP-121, Institute for Mathematics, Novosibirsk (unpublished).
[11] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1982a, Phys. Lett. **108B**, 134.
[12] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1982b, Z. Phys. C **16**, 55.
[13] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1984a, Sov. Phys. Usp. **27**, 161 [Usp. Fiz. Nauk **142**, 361 (1984)].
[14] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1984, Z. Phys. C **22**, 53.
[15] Achasov, N.N., S.A. Devyanin, and G.N. Shestakov, 1985, Z. Phys. C **27**, 99.
[16] Achasov, N.N., and G.N. Shestakov, 1988, Z. Phys. C **41**, 309.
[17] Achasov, N.N., and V.N. Ivanchenko, 1989, Nucl. Phys. B **315**, 465.

[18] Achasov, N.N., 1990, in *Proceedings of the Rheinfels Workshop 1990 on Hadron Mass Spectrum*, edited by E. Klempt and K. Peters, Nucl. Phys. B Proc. Suppl. **21**, 189 (1991).

[19] Achasov, N.N., and G.N. Shestakov, 1991, Sov. Phys. Usp. **34**, 471 [Usp. Fiz. Nauk **161**, 53 (1991)].

[20] Achasov, N.N., and G.N. Shestakov, 1992, Sov. J. Nucl. Phys. **55**, 1677 [Yad. Fiz. **55**, 2999 (1992)].

[21] Achasov, N.N., and G.N. Shestakov, 1994a, Phys. Rev. D **49**, 5779.

[22] Achasov, N.N., and G.N. Shestakov, 1994b, Mod. Phys. Lett. A **9**, 1351.

[23] Achasov, N.N., 1995, in *The Second DAΦNE Physics Handbook*, edited by L. Maiani, G. Pancheri, and N. Paver, Vol. II, p. 671.

[24] Achasov, N.N., V.V. Gubin, and E.P. Solodov, 1997, Phys. Rev. D **55**, 2672.

[25] Achasov, N.N., V.V. Gubin, and V.I. Shevchenko, 1997, Phys. Rev. D **56**, 203.

[26] Achasov, N.N., and V.V. Gubin, 1997, Phys. Rev. D **56**, 4084.

[27] Achasov, N.N., and V.V. Gubin, 1998, Phys. Rev. D **57**, 1987.

[28] Achasov, N.N., 1998, Physics-Uspekhi **41**, 1149 [Usp. Fiz. Nauk **168**, 1257 (1998)].

[29] Achasov, M.N., *et al.* (SND), 1998a, Phys. Lett. B **438**, 441.

[30] Achasov, M.N., *et al.* (SND), 1998b, Phys. Lett. B **440**, 442.

[31] Achasov, M.N., *et al.* (SND), 2000a, Phys. Lett. B **479**, 53.

[32] Achasov, M.N., *et al.* (SND), 2000b, Phys. Lett. B **485**, 349.

[33] Achasov, N.N., and G.N. Shestakov, 1999, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by G.V. Fedotovich and S.I. Redin (BINP, Novosibirsk, 2000), p. 294.

[34] Achasov, N.N., 1999, in *Proceedings of the 8th International Conference On Hadron Spectroscopy*, edited by W.G. Li, Y.Z. Huang, and B.S. Zou, Nucl. Phys. A **675**, 279c (2000).

[35] Achasov, N.N. and V.V. Gubin, 2001, Phys. Rev. D **63**, 094007.

[36] Achasov, N.N., 2001, in *Proceedings of the Ninth International Conference on Hadron Spectroscopy*, edited by D. Amelin and A.M. Zaitsev, AIP Conf. Proc. Vol. 619, p. 112 (2002).

[37] Achasov, N.N., 2002, Phys. At. Nucl. **65**, 546 [Yad. Fiz. **65**, 573 (2002)].

[38] Achasov, N.N., 2003a, Nucl. Phys. A **728**, 425.

[39] Achasov, N.N., 2003b, in *Proceedings of the International Symposium on Hadron Spectroscopy, Chiral Symmetry and Relativistic Description of Bound Systems*, edited by S. Ishida, K. Takamatsu, T. Tsuru, S.Y. Tsai, M. Ishida, and T. Komada, (KEK Proceedings 2003-7), p. 151.

[40] Achasov, N.N., 2003c, in *Proceedings of the KEK Workshop on Hadron Spectroscopy and Chiral Particle Search in J/Psi Decay Data at BES*, edited by K. Takamatsu, T. Tsuru, K. Ukai, and Z. Zheng (KEK Proceedings 2003-10), p. 66.

[41] Achasov, N.N. and G.N. Shestakov, 2003, Phys. Rev. D **67**, 114018.

[42] Achasov, N.N. and A.V. Kiselev, 2003, Phys. Rev. D **68**, 014006.

[43] Achasov, N.N., 2004a, in *Proceedings of the 13th International Seminar QUARKS'2004*, edited by D.G. Levkov, V.A. Matveev, and V.A. Rubakov (INR RAS, Moscow, 2004), p. 110.

[44] Achasov, N.N., 2004b, in *Proceedings of the International Bogolyubov Conference "Problems of Theoretical and Mathematical Physics"*, edited by V.G. Kadyshevsky and A.N. Sissakian, Phys. Part. Nucl. **36**, Suppl. 2, 146 (2005).

[45] Achasov, N.N., and G.N. Shestakov, 2004a, Phys. Rev. Lett. **92**, 182001.

[46] Achasov, N.N., and G.N. Shestakov, 2004b, Phys. Rev. D **70**, 074015.

[47] Achasov, N.N., and A.V. Kiselev, 2004, Phys. Rev. D **70**, 111901.

[48] Achasov, N.N., and G.N. Shestakov, 2005, Phys. Rev. D **72**, 013006.

[49] Achasov, N.N. and A.V. Kiselev, 2006, Phys. Rev. D **73**, 054029

[50] Achasov, N.N., A.V. Kiselev, and G.N. Shestakov, 2006, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by A. Bondar and S. Eidelman, Nucl. Phys. B Proc. Suppl. **162**, 127 (2006).

[51] Achasov, N.N., 2006, in *Proceedings of the 14th International Seminar QUARKS'2006*, edited by S.V. Demidov, V.A. Matveev, V.A. Rubakov, and G.I. Rubtsov (INR RAS, Moscow, 2007), Vol. 1, p. 29.

[52] Achasov, N.N. and A.V. Kiselev, 2007a, Phys. At. Nucl. **70**, 1956 [Yad.Fiz. **70**, 2005 (2007)].

[53] Achasov, N.N., A.V. Kiselev, 2007b, Phys. Rev. D **76**, 077501.

[54] Achasov, N.N., and G.N. Shestakov, 2007, Phys. Rev. Lett. **99**, 072001.

[55] Achasov, N.N., and G.N. Shestakov, 2008a, Phys. Rev. D **77**, 074020.

[56] Achasov, N.N., and G.N. Shestakov, 2008b, JETP Lett. **88**, 295 [Pisma Zh. Eksp. Teor. Fiz. **88**, 345(2008)].

[57] Achasov, N.N., A.V. Kiselev, and G.N. Shestakov, 2008, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by C. Bibi and G. Venanzoni, Nucl. Phys. B Proc. Suppl. **181-182**, 169 (2008).

[58] Achasov, N.N., A.V. Kiselev, 2008, Phys. Rev. D **78**, 058502.

[59] Achasov, N.N., 2008a, Invited Talk at *15th International Seminar on High Energy Physics: Quarks - 2008* (Sergiev Posad, Russia); eprint arXiv: 0810.2601 [hep-ph].

[60] Achasov, N.N., 2008b, in *Proceedings of the 14th High-Energy Physics International Conference on Quantum ChromoDynamics*, edited by S. Narison, Nucl. Phys. B Proc. Suppl. **186**, 283 (2009).

[61] Adachi I., *et al.* (Belle), 2008, Contributed to *the 34th International Conference on High Energy Physics* (Philadelphia, Pennsylvania), eprint arXiv: 0810.0334 [hep-ex].

[62] Adler, S.L., 1969, Phys. Rev. **177**, 2426.

[63] Akhmetshin, R.R., *et al.*(CMD-2), 1999a, Phys. Lett. B **462**, 371.

[64] Akhmetshin, R.R., *et al.*(CMD-2), 1999b, Phys. Lett. B **462**, 380.

[65] Albrecht, H., *et al.* (ARGUS), 1990, Z. Phys. C **48**, 183.

[66] Alde, D., *et al.* (GAMS), 1995, Z. Phys. C **66**, 375.

[67] Alde, D., *et al.* (GAMS), 1998, Eur. Phys. J. A **3**, 361.

[68] Alde, D., *et al.* (GAMS), 1999, Phys. At. Nucl. **62**, 421 [Yad. Fiz. **62**, 462 (1999)].

[69] Alford, M., and R.L. Jaffe, 2003, in *Proceedings of the High-Energy Physics Workshop on Scalar Mesons: An Interesting Puzzle for QCD*, edited by A.H. Fariborz, AIP Conf. Proc. Vol. 688, p. 208 (2003).

[70] Aloisio, A., *et al.* (KLOE), 2002a, Phys. Lett. B **536**, 209.

[71] Aloisio, A., *et al.* (KLOE), 2002b, Phys. Lett. B **537**, 21.

[72] Althoff, M., *et al.* (TASSO), 1985, Z. Phys. C **29**, 189.

[73] Althoff, M., *et al.* (TASSO), 1983, Phys. Lett. B **121**, 216.

[74] Ambrosino, F., *et al.* (KLOE), 2006, Phys. Lett. B **634** 148.

[75] Ambrosino, F., *et al.* (KLOE), 2007a, Eur. Phys. J. C **49** 473.

[76] Ambrosino, F., *et al.* (KLOE), 2007b, contributed to *23rd International Symposium on Lepton-Photon Interactions at High Energy* (Daegu, Korea); eprint arXiv: 0707.4609 [hep-ex].

[77] Ambrosino, F., *et al.* (KLOE), 2009a, eprint arXiv: 0903.4115 [hep-ex].

[78] Ambrosino, F., *et al.* (KLOE), 2009b, eprint arXiv: 0904.2539 [hep-ex].

[79] Amsler, C., and N.A. Törnqvist, 2004, Phys. Rep. **389**, 61.

[80] Amsler, C., *et al.* (Particle Data Group), 2008, Phys. Lett. B **667**, 1.

[81] Antreasyan, D., *et al.* (Crystal Ball), 1986, Phys. Rev. D **33**, 1847.

[82] Babcock, J., and J.L. Rosner, 1976, Phys. Rev. D **14**, 1286.

[83] Barash-Schmidt, N., *et al.* (Particle Data Group), 1969, Rev. Mod. Phys. **41**, 109.

[84] Barbieri, R., R. Gatto, and R. Kögerler, 1976, Phys. Lett. **60B**, 183.

[85] Bardeen, W.A., H. Fritzsch, and M. Gell-Mann, 1972, in *Proceedings of the Meeting on Scale and Conformal Symmetry in Hadron Physics*, edited by R. Gatto (Wiley, 1973); CERN-TH-1538; eprint arXiv: hep-ph/0211388.

[86] Barnes, T., 1985, Phys. Lett. B **165**, 434.

[87] Barnes, T., 1992, in *Proceedings of the 9th International Workshop on Photon-Photon Collisions*, edited by D.O. Caldwell and H.P. Paar (World Scientific, Singapore), p. 263, p. 275.

[88] Barrelet, E., 1972, Nuovo Cim. A **8**, 331.

[89] Behrend, H.J., *et al.* (CELLO), 1989, Z. Phys. C **43**, 91.

[90] Behrend, H.J., *et al.* (CELLO), 1992, Z. Phys. C **56**, 381.

[91] Bell, J.S., and L. Jackiw, 1969, Nuovo Cim. A **60**, 47.

[92] Berger, S.B., and B.T. Feld, 1973, Phys. Rev. D **8**, 3875.

[93] Berger, Ch., 1980, in *Proceedings of the International Workshop on gamma gamma Collisions* edited by G. Cochard and P. Kessler (Springer Verlag), Lecture Notes in Physics Vol. 134, p. 82.

[94] Berger, Ch., and W. Wagner, 1987, Phys. Rep. C **146**, 1.

[95] Berger, Ch., *et al.* (PLUTO), 1988, Z. Phys. C **37**, 329.

[96] Bernstein, A.M., 2007, eprint arXiv: 0707.4250 [hep-ph].

[97] Bienlein, J.K., 1992, in *Proceedings of the 9th International Workshop on Photon-Photon Collisions*, edited by D. Caldwell and H.P. Paar (World Scientific, Singapore), p. 241.

[98] Bijnens, J., and F. Cornet, 1988, Nucl. Phys. B **296**, 557.

[99] Bini, C. (KLOE), 2008, Talk given at *34th International Conference on High Energy Physics* (Philadelphia, Pennsylvania); eprint arXiv: 0809.5004 [hep-ex].

[100] Black, D., A.H. Fariborz, F. Sannino, and J. Schechter, 1998, Phys. Rev. **58**, 054012.

[101] Black, D., A.H. Fariborz, F. Sannino, and J. Schechter, 1999, Phys. Rev. **59**, 074026.

[102] Boglione, M., and M.R. Pennington, 1999, Eur. Phys. J. C **9**, 11.

[103] Bonvicini, G., *et al.* (CLEO), 2007, Phys. Rev. D **76**, 012001.

[104] Bossi, F., E. De Lucia, J. Lee-Franzini, S. Miscetti, and M. Palutan , 2008, Riv. Nuovo Cim. **031**, 531-623; eprint arXiv: 08111929 [hep-ex].

[105] Boyer, J., *et al.* (Crystal Ball), 1990, Phys. Rev. D **42**, 1350.

[106] Braccini, S., 2000, in *Proceedings of the MESON 2000 Workshop*, edited by L. Jarczyk, C. Guaraldo, H. Machner, and A. Magiera, Acta Phys. Polon. B **31**, 2143.

[107] Bramon, A., A. Grau, and G. Pancheri, 1992, Phys. Lett. B **289**, 97.

[108] Branz, T., T. Gutsche, and V.E. Lyubovitskij, 2007, Eur. Phys. J. A **37**, 303.

[110] Budnev, V.M., and A.E. Kaloshin, 1979, Phys. Lett. B **86**, 351.

[110] Bugg, D.V. ,2006, Eur. Phys. J. C **47**, 57.

[111] Bystritsky, Yu.M., M.K. Volkov, E.A. Kuraev, E. Bartoš, and M. Sečanský, 2007, Phys. Rev. D **77**, 054008.

[112] Cahn, R.N., 1989, in *Proceedings of the 14th International Symposium on Lepton and Photon at High Energies*, edited by M. Riordan (World Scientific, Singapore, 1990), p. 60.

[113] Caprini, I., G. Colangelo, and H. Leutwyler, 2006, Phys. Rev. Lett. **96**, 132001.

[114] Cavoto, G., 2007. in *Proceedings of the 5th Flavor Physics and CP Violation Conference* (Bled, Slovenia), p. 22; eprint arXiv:0707.1242 [hep-ex].

[115] Chanowitz, M.S., 1988, in *Proceedings of the 8th International Workshop on Photon-Photon Collisions*, edited by U. Karshon (World Scientific, Singapore) p. 205.

[116] Close, F.E., N. Isgur, and S. Kumano, 1993, Nucl. Phys. B **389**, 513.

[117] Close, F.E., and N.A. Törnqvist, 2002, J. Phys. G **28**, R249.

[118] Cordier, A., 1984, *Proceedings of the 6th International Workshop on Photon-Photon Collisions*, edited by R.L. Lander, (World Scientific, Singapore, 1985), p. 122.

[119] Delbourgo, R., M.D. Scadron, 1998, Int. J. Mod. Phys. A **13**, 657.

[120] Di Micco, B., 2008, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by C. Bibi and G. Venanzoni, Nucl. Phys. B Proc. Suppl. **181-182**, 215 (2008).

[121] Donoghue, J.F., B.R. Holstein, and Y.C. Lin, 1988, Phys. Rev. D **37**, 2423.

[122] Donoghue, J.F., and B.R. Holstein, 1993, Phys. Rev. D **48**, 137.

[123] Dorofeev, V., *et al.*, 2007, in *Proceedings of the 12th International Conference on Hadron Spectroscopy* (Fras-

cati Physics Series), Vol. XLVI; eprint arXiv: 0712.2512 [hep-ex].

[124] Dorofeev, V., *et al.*, 2008, Eur. Phys. J. A **38**, 149.

[125] Dubrovin, M. (for the CLEO Collaboration), 2003, in *High-Energy Physics Workshop on Scalar Mesons: An Interesting Puzzle for QCD*, edited by A. H. Fariborz, AIP Conf. Proc. Vol. 688, p. 231 (2004).

[126] Durusoy, N.B., *et al.*, 1973, Phys. Lett. B **45**, 517.

[127] Dzierba, A.R., 1995, in *Proceedings of the Second Workshop on Physics and Detectors for DAΦNE'95*, edited by R. Baldini, F. Bossi, G. Capon, and G. Pancheri (Frascati Physics Series, 1996), Vol. 4, p. 99.

[128] Ebert, D., R.N. Faustov, and V.O. Galkin, 2009, Eur. Phys. J. C **60**, 273.

[129] Edwards, C., *et al.* (Crystal Ball), 1982, Phys. Lett. B **110**, 82.

[130] Erne, F.C., 1984, *Proceedings of the 6th International Workshop on Photon-Photon Collisions*, edited by R.L. Lander (World Scientific, Singapore, 1985), p. 151.

[131] Estabrooks, P., 1979, Phys. Rev. D **19**, 2678.

[132] Faiman, D., H.J. Lipkin, and H.R. Rubinstein, 1975, Phys. Lett. B **59**, 269.

[133] Fariborz, A.H., R. Jora, and J. Schechter, 2006, Phys. Rev. D **76**, 014011.

[134] Fariborz, A.H., R. Jora, and J. Schechter, 2008a, Phys. Rev. D **77**, 094004.

[135] Fariborz, A.H., R. Jora, and J. Schechter, 2008b, eprint arXiv: 0810.4640 [hep-ph].

[136] Fariborz, A.H., R. Jora, and J. Schechter, 2009, eprint arXiv: 0902.2825 [hep-ph].

[137] Feindt, M., and J. Harjes, 1990, in *Proceedings of the Rheinfeld 1990 Workshop on the Hadron Mass Specrum*, edited by E. Klempt and K. Peters, Nucl. Phys. B Proc. Suppl. **21**, 61 (1991).

[138] Feldmann, T., 2000, Int. J. Mod. Phys. A **15**, 159.

[139] Field, J.H., 1981, in *Proceedings of the 4th International Colloquium on Photon-Photon Interactions* (World Scientific, Singapore), p. 447.

[140] Flatte, S.M., *et al.*, 1972, Phys. Lett. B **38**, 232.

[141] Flatte, S.M., 1976, Phys. Lett. B **63**, 224, 228.

[142] Gay, J., *et al.*, 1976, Phys. Lett. B **63**, 220.

[143] Gell-Mann, M., and M.L. Goldberger, 1954, Phys. Rev. **96**, 1433.

[144] Gell-Mann, M., and M. Levy, 1960, Nuovo Cim. **16**, 705.

[145] Gell-Mann, M., 1964, Physics **1**, 63.

[146] Gersten, A., 1969, Nucl. Phys. B **12**, 537.

[147] Godfrey, S., and J. Napolitano, 1999, Rev. Mod. Phys. **71** 1411.

[148] Grayer, G., *et al.*, 1974, Nucl. Phys. **B75**, 189.

[149] Groom, D.E., *et al.* (Particle Data Group), Eur. Phys. J. C **15**, 1.

[150] Gunter, J., *et al.* (E852), 2001, Phys. Rev. D **64**, 072003.

[151] Hikasa, K., *et al.* (Particle Data Group), 1992, Phys. Rev. D **45**, S1.

[152] Hilger, E., 1981, in *Proceedings of the 4th International Colloquium on Photon-Photon Interactions* (World Scientific, Singapore), p. 149.

[153] Hoogland, W., *et al.*, 1977, Nucl. Phys. B **126**, 109.

[154] Hyams, B. *et al.*, 1973, Nucl. Phys. B **64**, 134.

[155] Hyams, B. *et al.*, 1975, Nucl. Phys. B **100**, 205.

[156] Ioffe, B.L., A.G. Oganesian, 2007, Phys. Lett. B **647**, 389.

[157] Ishida, S., M.Y. Ishida, H. Takahashi, T. Ishida, K. Takamatsu, and T. Tsuru, 1996, Prog. Theor. Phys. **95**, 745.

[158] Ishida, S., 1997, in *Proceedings of the 7th International Conference on Hadron Spectroscopy*, edited by S.-U. Chung and H. Willutzki, AIP Conf. Proc. Vol. 432, p. 705 (1998).

[159] Ishida, M., 2000, in *Proceedings of the Possible Existence of sigma-Meson and Its Implication to Hadron Physics*, edited by S. Ishida, M.Y. Ishida, T. Kunihiro, T. Hatsuda, M. Oka, H. Shimizu, K. Takamatsu, T. Tsuru, and C.S. Lim (KEK Proceedings 2000-4); eprint arXiv: hep-ph/0012325.

[160] Ivashin, S., and A. Korchin, 2008, Eur. Phys. J. C **54**, 89.

[161] Jackson, J.D., 1976, in *Proceedings of the SLAC Summer Institute on Particle Physics: Weak Interactions at High Energy and the Production of New Particles*, edited by M.C. Zipf, SLAC Rep. No. 198, p. 147.

[162] Jaffe, R.L., 1977, Phys. Rev. D **15**, 267, 281.

[163] Jaffe, R.L., and F. Wilczek, 2003, Phys. Rev. Lett. **91**, 232003.

[164] Jaffe, R.L., 2005, Phys. Rep. **409**, 1 [Nucl. Phys. B Proc. Suppl. **142**, 343 (2005)].

[165] Jaffe, R.L., 2006, in *Proceedings of the YKIS Seminar on New Frontiers in QCD: Exotic Hadrons and Hadronic Matter* (Kyoto, Japan), Prog. Theor. Phys. Suppl. **168**, 127 (2007) [AIP Conf. Proc. Vol. 964, p. 1 (2007)].

[166] Johnson, R.P., 1986, Ph.D. thesis, SLAC-Report-294.

[167] Kalashnikova, Yu.S., A.E. Kudryavtsev, A.V. Nefediev, C. Hanhart, and J. Haidenbauer, 2005, Eur. Phys. J. A **24**, 437.

[168] Kalinovsky, Yu.L., and M.K. Volkov, 2008, eprint arXiv: 0809.1795 [hep-ph].

[169] Kaloshin, A.E., and V.V. Serebryakov, 1986, Z. Phys. C **32**, 279.

[170] Karch, K.-H., 1991, in *Proceedings of the 26th Rencontre de Moriond: High Energy Hadronic Interactions*, edited by J. Tran Thanh Van (Gif-sur-Yvette, France, Editions Frontieres), p. 423.

[171] Klempt, E., and A. Zaitsev, 2007, Phys. Rep. **454**, 1.

[172] Kolanoski, H., 1984, in *Springer Tracts in Modern Physics* (Springer-Verlag, Berlin), Vol. 105, p. 1.

[173] Kolanoski, H., 1985, in *Proceedings of the 12th International Symposium on Lepton and Photon Interactions at High Energies*, edited by M. Konuma and K. Takahashi (Kyoto Univ., Research Inst. Fund. Phys., 1986), p. 90.

[174] Kolanoski, H., and P. Zerwas, 1987, in *High Energy Electron-Positron Physics*, edited by A. Ali and P. Söding (World Scientific, Singapore, 1988), p. 695.

[175] Kolanoski, H., 1988, in *Proceedings of the 9th European Symposium on Antiproton-Proton Interactions and Fundamental Symmetries*, edited by K. Kleinknecht and E. Klempt, Nucl. Phys. B Proc. Suppl. **8**, 41 (1989).

[176] Kolanoski, H., 1991 in *Proceedings of the 4th International Conference on Hadron Spectroscopy*, edited by S. Oneda and D.C. Peaslee (World Scientific, Singapore, 1992), p. 377.

[177] Krammer, M., and H. Krasemann, 1978, Phys. Lett. B **73**, 58.

[178] Krammer, M., 1978, Phys. Lett. B **74**, 361.

[179] Krasemann, H., and J.A.M. Vermaseren, 1981, Nucl. Phys. B **184**, 269.

[180] Lasinski, T.A., *at al.* (Particle Data Group), 1973, Rev. Mod. Phys. **45**, S1.

[181] Leutwyler, H., 1998, Nucl. Phys. B Proc. Suppl. **64**, 223.

[182] Leutwyler, H., 2008, in *Proceedings of the Workshop on Scalar Mesons and Related Topics Honoring 70th Birthday of Michael Scadron*, AIP Conf. Proc. Vol. 1030, p. 46.

[183] Levy, M., 1967, Nuovo Cim. A **52**, 23.

[184] Li, Z.P., F.E. Close, and T. Barnes, 1991, Phys. Rev. D **43**, 2161.

[185] Li, D-M., H. Yu, and Q-X. Shen, 2001, J. Phys. G **27**, 807.

[186] Low, F.E., 1954, Phys. Rev. **96**, 1428.

[187] Lucio, J.L., and M. Napsuciale, 1994, Phys. Lett. B **331**, 418.

[188] Lyth, D.H., 1984, J. Phys. G **10**, 39.

[189] Lyth, D.H., 1985, J. Phys. G **11**, 459.

[190] Maiani, L., A. Polosa, F. Piccinini, and V. Riquer, 2004, Phys. Rev. Lett. **93**, 212002.

[191] Maiani, L., A.D. Polosa, and V. Riquer, 2007, Phys. Lett. B **651**, 129.

[192] Mao, Y., X.-G. Wang, O. Zhang, H. Q. Zheng, and Z. Y. Zhou, 2009, eprint arXiv: 0904.1445 [hep-ph].

[193] Marsiske, H., *et al.* (MARK II), 1990, Phys. Rev. D **41**, 3324.

[194] Martin, A.D., E.N. Ozmutlu, and E.J. Squires, 1977, Nucl. Phys. B **121**, 514.

[195] Mennessier, G., 1983, Z. Phys. C **16**, 241.

[196] Mennessier, G., P. Minkowski, S. Narison, and W. Ochs, 2007, eprint arXiv: 0707.451 [hep-ph].

[197] Mennessier, G., S. Narison, and W. Ochs, 2008a, Phys. Lett. B **665**, 205.

[198] Mennessier, G., S. Narison, and W. Ochs, 2008b, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by C. Bibi and G. Venanzoni, Nucl. Phys. B Proc. Suppl. **181-182**, 238 (2008).

[199] Mennessier, G., 2008, Talk given at *the 14th High-Energy Physics International Conference in Quantum Chromodynamics* (Montpellier, France); eprint arXiv: 0811.1589 [hep-ph].

[200] Montanet, L., 1983, Rept. Prog. Phys. **46**, 337.

[201] Morgan, D., 1974, Phys. Lett. B **51**, 71.

[202] Morgan, D., and M.R. Pennington, 1987, Phys. Lett. B **192**, 207.

[203] Morgan, D., and M.R. Pennington, 1988, Z. Phys. C **37**, 431.

[204] Morgan, D., and M.R. Pennington, 1990, Z. Phys. C **48**, 623.

[205] Morgan, D., and M.R. Pennington, 1991, Phys. Lett. B **272**, 134.

[206] Morgan, D., M.R. Pennington, and M.R. Whalley, 1994, *A Compilation of the Data on Two-Photon Reactions Leading to Hadron Final States*, J. Phys. G **20**, Suppl. 8A, A1.

[207] Mori, T., *et al.* (Belle), 2003, in *Proceedings of the International Simposium on Hadron Spectroscopy, Chiral Symmetry and Relativistic Description of Bound Systems*, edited by S. Ishida, K. Takamatsu, T. Tsuru, S.Y. Tsai, M. Ishida, and T. Komada (KEK Proceedings 2003-7), p. 159.

[208] Mori, T., *et al.* (Belle), 2007a, Phys. Rev. D **75**, 051101(R).

[209] Mori, T., *et al.* (Belle), 2007b, J. Phys. Soc. Jpn. **76**, 074102.

[210] Münz, C.R., 1996, Nucl. Phys. A **609**, 364.

[211] Nakazawa, N., 2008, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by C. Bibi and G. Venanzoni, Nucl. Phys. B Proc. Suppl. **181-182**, 233 (2008).

[212] Narison, S., 2006, Phys. Rev. D **73**, 114024.

[213] Narison, S., 2008, Talk given at *the 14th High-Energy Physics International Conference in Quantum Chromodynamics* (Montpellier, France); eprint arXiv: 0811.0563 [hep-ph].

[214] Oest, T., *et al.* (JADE), 1990, Z. Phys. C **47**, 343.

[215] Oller, J.A., and E. Oset, 1997, in *Proceedings of the 7th International Conference on Hadron Spectroscopy*, edited by S.-U. Chung and H. Willutzki, AIP Conf. Proc. Vol. 432, p. 413 (1998).

[216] Oller, J.A., L. Roca, and C. Schat, 2008, Phys. Lett. B **659**, 201.

[217] Oller, J.A., and L. Roca, 2008, Eur. Phys. J. A **37**, 15.

[218] Olsson, J.E., 1983, in *Proceedings of the 5th International Workshop on Photon-Photon Interactions*, edited by C. Berger, Lecture Notes in Physics (Springer-Verlag, Berlin), Vol. 191, p. 45.

[219] Pennington, M.R., 1999, in *Proceedings of the International Conference on the Structure and Interactions of the Photon*, edited by S. Soldner-Rembold, Nucl. Phys. B Proc. Suppl. **82**, 291 (2000).

[220] Pennington, M.R., 2006, Phys. Rev. Lett. **97**, 011601.

[221] Pennington, M.R., 2007a, Progr. Theor. Phys. Suppl. **168**, 143.

[222] Pennington, M.R., 2007b, in *Proceedings of the 11th International Conference on Meson-Nucleon Physics and the Structure of the Nucleon* (Julich, Germany), p. 106.

[223] Pennington, M.R., 2007c, Mod. Phys. Lett. A **22**, 1439.

[224] Pennington, M.R., 2008, in *Proceedings of the International Workshop on e^+e^- Collisions from ϕ to ψ*, edited by C. Bini and G. Venanzoni, Nucl. Phys. B Proc. Suppl. **181-182**, 251 (2008).

[225] Pennington, M.R., T. Mori, S. Uehara, and Y. Watanabe, 2008, Eur. Phys. J. C **56**, 1.

[226] Petersen, J.L., 1977, Yellow CERN Preprint 77-04.

[227] Poppe, M., 1986, Intern. J. Mod. Phys. A **1**, 545.

[228] Protopopescu, S.D., *et al.*, 1973, Phys. Rev. D **7**, 1279.

[229] Rittenberg, A., *at al.* (Particle Data Group), 1971, Rev. Mod. Phys. **43**, S1.

[230] Rosenfeld, A.H., *et al.* (Particle Data Group), 1965, Rev. Mod. Phys. **37**, 633.

[231] Rosenfeld, A.H., *et al.* (Particle Data Group), 1967, Rev. Mod. Phys. **39**, 1.

[232] Rosner, J.L., 1981, Phys. Rev. D **23**, 1127.

[233] Sadovsky, S.A., 1999, Phys. Atom. Nucl. **62**, 519 [Yad. Fiz. **62**, 562 (1999)].

[234] Sannino, F., and J. Schechter, 1995, Phys. Rev. D **52**, 96.

[235] Shekhovtsova, O., G. Venanzoni, and G. Pancheri, 2009, eprint arXiv:0901.4440 [hep-ph].

[236] Spanier, S., N.A. Törnqvist, and C. Amsler, 2008, Phys. Lett. B **667**, 594.

[237] 't Hooft, G., G. Isidori, L. Maiani, A.D. Polosa, and V. Riquer, 2008, Phys. Lett. B **662**, 424.

[238] Törnqvist, N.A., 1995, Z. Phys. C **68**, 647.

[239] Törnqvist, N.A., 2007, Acta. Phys. Polon. B **38**, 2831.

[240] Tuan, S.F., 2001, in *Proceedings of the Ninth International Conference on Hadron Spectroscopy*, edited by D. Amelin and A.M. Zaitsev, AIP Conf. Proc. Vol. 619, p. 495 (2002).

[241] Tuan, S.F., 2003, in *Proceedings of the International Symposium on Hadron Spectroscopy, Chiral Symmetry and Relativistic Description of Bound Systems*, edited by S. Ishida, K. Takamatsu, T. Tsuru, S.Y. Tsai, M. Ishida, and T. Komada, (KEK Proceedings 2003-7), p. 319.

[242] Uehara, S., *et al.* (Belle), 2008, Phys. Rev. D **78**, 052004.

[243] van Beveren, E., and G. Rupp, 2007, in *Proceedings of 11th International Conference on Meson-Nucleon Physics and the Structure of the Nucleon*, edited by H. Machner and S. Krewald, p. 130.

[244] van Beveren, E., F. Kleefeld, and G. Rupp, 2008, eprint arXiv: 0811.2589 [hep-ph].

[245] Watson, K.M., 1952, Phys. Rev. **88**, 1163.

[246] Wedemeyer, R.J., 1981, in *Proceedings of the 10th International Symposium on Lepton and Photon Interactions at High Energies*, edited by W. Pfeil (Bonn Univ., Phys. Inst.), p. 410.

[248] Weinstein, J., and N. Isgur, 1982, Phys. Rev. Lett. **48**, 659.

[248] Weinstein, J., and N. Isgur, 1990, Phys. Rev. D **41**, 2236.

[249] Wu, J.-J., Q. Zhao, and B.S. Zou, 2007, Phys. Rev. D **75**, 114012.

[250] Wu, J.-J., and B.S. Zou, 2008, Phys. Rev. D **78**, 074017.

[251] Yao, W.-M., *et al.* (Particle Data Group), 2006, J. Phys. G **33**, 1.